

Overall summary


- All indicators of influenza activity remain exceptionally low for this time of year, despite continued monitoring for influenza across Canada.
- To date this season, there has been no evidence of community circulation of influenza despite continued testing above seasonal levels. Influenza activity has remained below the threshold required to declare the start of the 2020-21 influenza season.
- Three influenza-like-illness (ILI) outbreaks in schools and/or daycares were reported in weeks 10 to 11, for a total of 124 ILI outbreaks reported this season to date. No laboratory-confirmed outbreaks of influenza have been reported to date this season.
- 12,105 participants reported to FluWatchers and 19 (0.16%) participants reported cough and fever, which remains low compared to previous seasons. Provide valuable information to track flu and COVID-19 across Canada: [Sign up to become a FluWatcher!](#)
- The 2020-2021 Seasonal Influenza Immunization Coverage Survey showed that influenza vaccine coverage was similar to previous seasons.
- Influenza surveillance indicators are influenced by the COVID-19 pandemic, including changes in healthcare-seeking behaviour, impacts of public health measures and influenza testing practices. Current data should be interpreted with consideration to this context. See the [COVID-19 Epidemiology update](#) for information on COVID-19 cases in Canada.
- Have you already forgotten what happened last flu season? Not to worry, FluWatch has got you covered! Check out the [2019-2020 FluWatch annual report](#) for Canada's official analysis on the entire 2019-2020 influenza season.

Influenza/influenza-like illness (ILI) activity (geographic spread)

During week 11, no influenza/ILI activity was reported (Figure 1).

Figure 1 – Map of influenza/ILI activity by province and territory, Canada, week 2021-11

Number of Regions Reporting in Week 11: 53 out of 53



Laboratory-confirmed influenza detections

In weeks 10 to 11, two laboratory detections of influenza were reported.

To date this season, 66 influenza detections have been reported (Figure 2), which is significantly lower than the past six seasons where an average of 43,097 influenza detections were reported for the season to date. All provinces and territories are closely monitoring indicators of influenza activity this season. Data in the FluWatch report represent surveillance data available at the time of writing, and may change as updates are received.

Thirty-one of the influenza detections reported to date this season are known to be associated with recent live attenuated influenza vaccine (LAIV) receipt and do not represent community circulation of seasonal influenza viruses. LAIV strains are attenuated but can be recovered by nasal swab in children and adults following vaccination with that product (i.e., "shedding"). For more information, please refer to the [Canadian Immunization Guide Chapter on Influenza and Statement on Seasonal Influenza Vaccine for 2020–2021 \(Section IV.2\)](#).

Overall, the percentage of laboratory tests positive for influenza remains at exceptionally low levels, despite the elevated levels of testing seen this week. In weeks 10 to 11, 42,855 tests for influenza were performed at reporting laboratories and the average percentage of tests positive was 0%. Compared to the past six seasons, the number of tests performed for this time period was higher than the average (22,040) and the percentage of tests positive for influenza remains well below average (22.3%) (Figure 3a and 3b).


In Canada, at the national level, the influenza season typically starts around week 47 (mid-November) and is defined when at least 5% of influenza tests are positive and a minimum of 15 positive tests are observed nationally. Over the past 6 seasons, the influenza season has begun as early as week 43 (mid-October) and as late as week 1 (early January). As of week 11, influenza activity has remained below the threshold required to declare the start of the 2020-2021 influenza season.

Testing for influenza and other respiratory viruses has been influenced by the current COVID-19 pandemic. Changes in laboratory testing practices may affect the comparability of data to previous weeks or previous seasons.

For more detailed weekly and cumulative influenza data, see the text descriptions for [Figure 2](#) or the [Respiratory Virus Detections in Canada Report](#).

Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, weeks 2020-35 to 2021-11

Number of laboratories reporting in week 11: 31 out of 35




The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

Figure 3 – Number of influenza tests (a) and percentage of tests positive (b) in Canada by week, weeks 2020-35 to 2021-11 as compared to previous seasons

a) Current number of influenza tests performed, compared to previous seasons

b) Current percentage of tests positive for influenza, compared to previous seasons



The shaded area represents the maximum and minimum number of influenza tests (a) or percentage of tests positive (b) reported by week from seasons 2014-2015 to 2019-2020.

Syndromic / influenza-like illness (ILI) surveillance


Healthcare practitioners sentinel syndromic surveillance

In weeks 10 to 11, an average of 0.3% of visits to healthcare professionals were due to influenza-like illness (ILI), which remains lower than average compared to recent seasons (Figure 4).

This trend should be interpreted with caution as there have been changes in healthcare seeking behavior of individuals and a smaller number of sentinels reporting compared to previous seasons.

Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, weeks 2020-35 to 2021-11

Number of Sentinels Reporting in Week 11: 50



The shaded area represents the maximum and minimum percentage of visits for ILI reported by week from seasons 2014-2015 to 2019-2020.


FluWatchers

In week 11, 12,105 participants reported to FluWatchers, and a total of 19 participants (0.16%) reported symptoms of cough and fever (Figure 5). The percentage of participants reporting cough and fever remains at very low levels and may be a direct effect of individual and public health measures enacted to reduce the spread of COVID-19. FluWatchers reporting is not impacted by changes in health services or health seeking behaviours.

If you are interested in becoming a [FluWatcher](#), [sign up today](#).

Figure 5 – Percentage of FluWatchers participants reporting cough and fever, Canada, weeks 2020-35 to 2021-11


Number of Participants Reporting in Week 11: 12,105



The shaded area represents the maximum and minimum percentage of percentage of participants reporting cough and fever by week, from seasons 2014-2015 to 2019-2020

Online Figure – Geographic distribution of FluWatchers participants reporting cough and fever, Canada, week 2021-11

Click on the map to access the link



Influenza outbreak surveillance

In weeks 10 to 11, three influenza-like-illness (ILI) outbreaks in schools and/or daycares were reported.

To date this season, 124 influenza-like-illness (ILI) outbreaks in schools and/or daycares, and no other influenza or ILI outbreaks have been reported in any other settings.

Outbreaks of ILI in schools and daycares are identified when greater than 10% absenteeism due to ILI are reported in these settings. The number of ILI outbreaks in schools and daycares reported is higher compared to the same time period in the previous two seasons. This is not unexpected given changes to outbreak surveillance, specifically increased vigilance in schools to monitor and report absenteeism due to ILI, and the increased restrictions on attendance for children with symptoms of viral respiratory illness.

Outbreaks of ILI are not specific to any one respiratory pathogen and can be due influenza or other respiratory viruses, including rhinovirus and SARS-CoV-2, the virus that causes COVID-19. For more information on the respiratory viruses currently circulating in Canada, please refer to the [Respiratory Virus Detections in Canada Report](#).

Number of provinces and territories reporting in weeks 10 to 11¹: 13 out of 13

¹All Provinces and Territories (PTs) participate in FluWatch's outbreak surveillance system. This outbreak system monitors influenza and ILI outbreaks in long-term care facilities, acute care facilities, schools and daycares, remote and/or isolated communities, and facilities categorized as 'other'. Not all reporting PTs report outbreaks in all these settings. All PTs report laboratory confirmed outbreaks in LTCF. Four PTs (NB, NL, NS and YK) report ILI outbreaks in schools and/or daycares.

Severe outcomes influenza surveillance

Provincial/territorial influenza hospitalizations and deaths

In weeks 10 to 11, no influenza-associated hospitalizations were reported by participating provinces and territories². To date this season no influenza-associated hospitalizations were reported by participating provinces and territories.

Number of provinces and territories reporting in weeks 10 to 11: 9 out of 9

²Influenza-associated hospitalizations are reported by Alberta, Manitoba, New Brunswick, Newfoundland and Labrador, Northwest Territories, Nova Scotia, Prince Edward Island and Yukon. Only hospitalizations that require intensive medical care are reported by Saskatchewan.

Pediatric influenza hospitalization and deaths

In weeks 10 to 11 no influenza-associated hospitalizations were reported. To date this season, less than five pediatric (≤ 16 years of age) hospitalizations with influenza have been reported by the Immunization Monitoring Program Active (IMPACT) network.

Adult influenza hospitalization and deaths

Surveillance of laboratory-confirmed influenza-associated adult (≥ 16 years of age) hospitalizations by the Canadian Immunization Research Network (CIRN) Serious Outcomes Surveillance (SOS) network has not begun for the 2020-2021 season.

Influenza strain characterizations

Due to the very low influenza circulation to date this season, the National Microbiology Laboratory has not received influenza viruses collected during the 2020-2021 season for strain characterization.

Antiviral resistance

Due to the very low influenza circulation to date this season, the National Microbiology Laboratory has not received influenza viruses collected during the 2020-2021 season for antiviral resistance testing.

Vaccine monitoring

Vaccine monitoring refers to activities related to the monitoring of influenza vaccine coverage and effectiveness.

Vaccine coverage

The Seasonal Influenza Immunization Coverage Survey is an annual telephone survey conducted between January and February that collects information from Canadians on whether they received the annual seasonal influenza vaccine that season. Vaccine coverage is measured as the percentage of people who reported receiving the influenza vaccine in a specific influenza season.

In the 2020-21 influenza season, coverage was similar to the 2019-20 season, at:

- 32% among adults aged 18 to 64 years.
 - 29% among adults aged 18-64 without chronic medical conditions.
 - 41% among adults aged 18 to 64 years with chronic medical conditions.
- 70% among seniors (aged 65 years and older).

Table 1 – Influenza vaccine coverage among adults (n=3,032)* by risk group† and gender‡, Seasonal Influenza Vaccination Coverage Survey, Canada, September 2020 – February 2021

Age group (years)	All		Male		Female	
	N	Vaccine Coverage % (95% CI)	N	Vaccine Coverage % (95% CI)	N	Vaccine Coverage % (95% CI)
All adults (≥18)	3014	40.4 (38.4-42.4)	1399	35.2 (32.5-38.0)	1595	45.2 (42.4-48.1)
18-64	2152	32.4 (30.1-34.6)	1024	27.3 (24.3-30.3)	1115	37.4 (34.1-40.7)
with chronic medical conditions	646	40.5 (36.2-44.8)	286	36.9 (30.7-43.1)	354	43.5 (37.5-49.4)
without chronic medical conditions	1498	29.2 (26.6-31.8)	734	24.2 (20.8-27.6)	757	34.6 (30.6-38.5)
≥65	862	70.4 (67.1-73.8)	375	68.0 (62.8-73.2)	480	72.8 (68.4-77.1)

*18 people did not recall whether they had received the influenza vaccine and were excluded from coverage estimates.

†11 people who were 18-64 years old did not disclose whether they had any chronic medical conditions (CMC) and were excluded from stratified analysis.

‡11 people did not disclose their gender and 9 people did not identify themselves as male or female. They were excluded from stratified analysis.

Vaccine effectiveness

Influenza vaccine effectiveness estimates are typically available in February or March of each year. Given the low influenza circulation this season to date, VE estimates will not be available for the 2020-2021 season.

Provincial/territorial and international surveillance links

- British Columbia – [Influenza Surveillance; Vaccine Effectiveness Monitoring](#)
- Alberta – [Respiratory Virus Surveillance](#)
- Saskatchewan – [Influenza Reports](#)
- Manitoba – [Seasonal Influenza Reports](#)
- Ontario – [Ontario Respiratory Pathogen Bulletin](#)
- Québec – [Système de surveillance de la grippe](#) (available in French only)
- New Brunswick – [Influenza Surveillance Reports](#)
- Prince Edward Island – [Influenza Summary](#)
- Nova Scotia – [Respiratory Watch Report](#)
- Newfoundland and Labrador – [Surveillance and Disease Reports](#)
- Yukon – [Information on Pandemic, Influenza, Seasonal Flu, Avian Flu and H1N1](#)
- Northwest Territories – [Influenza/ Flu Information](#)
- Nunavut – [Influenza Information](#)
- World Health Organization - [Global Influenza Programme, Influenza Updates](#)
- Pan American Health Organization – [Influenza situation report](#)
- U.S. Centers for Disease Prevention & Control (CDC) - [Weekly Influenza Summary Update](#)
- ECDC – [Surveillance reports and disease data on seasonal influenza](#)
- United Kingdom – [Weekly Influenza Activity Reports](#)
- Hong Kong Centre for Health Protection - [Flu Express](#)
- Australia – [Influenza Surveillance Report and Activity Updates](#)
- New Zealand – [Influenza Weekly Update](#)

Notes

The data in the FluWatch report represent surveillance data available at the time of writing. All data are preliminary and may change as more reports are received.

To learn more about the FluWatch program, see the [Overview of influenza monitoring in Canada](#) page. For more information on the flu, see our [Flu \(influenza\)](#) web page.

THANK YOU to our Fluwatch surveillance partners who participate the influenza surveillance program!

This [report](#) is available on the Government of Canada Influenza webpage.

Ce [rapport](#) est disponible dans les deux langues officielles.