



Health  
Canada Santé  
Canada

Your health and  
safety... our priority.

Votre santé et votre  
sécurité... notre priorité.

*State of the Science Report  
for a Screening Health Assessment*

# **Perfluorooctane Sulfonate (PFOS)**

**Its Salts and Its Precursors  
that Contain the  
 $C_8F_{17}SO_2$  or  $C_8F_{17}SO_3$  Moiety**



Canada 

“Health Canada is the federal department responsible for helping Canadians maintain and improve their health. We assess the safety of drugs and many consumer products, help improve the safety of food, and provide information to Canadians to help them make healthy decisions. We provide health services to First Nations people and to Inuit communities. We work with the provinces to ensure our health care system serves the needs of Canadians.”

Published by authority of the  
Minister of Health

Également disponible en français sous le titre :  
*Rapport sur l'état des connaissances scientifiques sous-jacentes  
à une évaluation préalable des risques pour la santé  
Le sulfonate de perfluorooctane, ses sels et ses précurseurs  
contenant la fraction  $C_8F_{17}SO_2$  ou  $C_8F_{17}SO_3$*

This publication can be made available on request on diskette, large print,  
audio-cassette and braille.

*State of the Science Report  
for a Screening Health Assessment*

# **Perfluorooctane Sulfonate (PFOS)**

**Its Salts and Its Precursors  
that Contain the  
 $C_8F_{17}SO_2$  or  $C_8F_{17}SO_3$  Moiety**

## **Table of Contents**

Introduction ... **1**

Identity, Uses and Sources of Exposure ... **2**

Exposure Assessment and Hazard Characterization ... **4**


Conclusion for Human Health ... **6**

References ... **16**

Table 1: Summary of health effects information for PFOS and related compounds ... **9**

Appendix 1: PFOS and Related Substances

**Perfluorooctane Sulfonate, Its Salts and Its  
Precursors that Contain the C<sub>8</sub>F<sub>17</sub>SO<sub>2</sub> or  
C<sub>8</sub>F<sub>17</sub>SO<sub>3</sub> Moiety**



## Introduction

Under the *Canadian Environmental Protection Act, 1999* (CEPA 1999) the Minister of Health may gather information, conduct investigations and evaluations, including screening assessments, relevant for the purpose of assessing whether a substance is entering or may enter the environment in a quantity or concentration or under conditions that constitute or may constitute a danger in Canada to human life or health.

Screening health assessments focus initially on conservative assessment of hazard or effect levels for critical endpoints and upper-bounding estimates of exposure, after consideration of all relevant identified information. Decisions based on the nature of the critical effects and margins between conservative effect levels and estimates of exposure take into account confidence in the completeness of the identified databases on both exposure and effects, within a screening context. Additional background information on screening health assessments conducted under this program is available at [http://www.hc-sc.gc.ca/ewh-semt/contaminants/existsub/index\\_e.html](http://www.hc-sc.gc.ca/ewh-semt/contaminants/existsub/index_e.html).

A screening health assessment was undertaken on perfluorooctane sulfonate (PFOS), its salts and its precursors containing the C<sub>8</sub>F<sub>17</sub>SO<sub>2</sub> or C<sub>8</sub>F<sub>17</sub>SO<sub>3</sub> moiety on the basis that some of these compounds were included in the Domestic Substances List pilot phase for screening and in response to a request to the Minister of the Environment to add these compounds to the Priority Substances List.

The State of the Science Report for a screening assessment and associated unpublished supporting working documentation were prepared by evaluators within the Existing Substances Division of Health Canada; the content of these documents was reviewed at several meetings of senior Divisional staff. The draft Report was subsequently externally reviewed for adequacy of

data coverage and defensibility of the conclusions. The draft report was also reviewed during a 60-day public comment period. The supporting working documentation is available upon request by e-mail from [ExSD@hc-sc.gc.ca](mailto:ExSD@hc-sc.gc.ca)

Information identified as of September 2003 was considered for inclusion in this screening health assessment. Additional information identified after that time and its relevance to the scientific assessment is presented in the last section of this document. The critical information and considerations upon which the assessment is based are summarized below.

### **Identity, Uses and Sources of Exposure**

PFOS, its salts and its precursors form part of a larger chemical class of fluorochemicals typically referred to as perfluorinated alkyl compounds (PFAs). Depending upon the intended use, the various precursors of PFOS are formed via derivatization of perfluorooctanesulfonylfluoride (POSF:  $C_8F_{17}SO_2F$ ) (OECD, 2002), yielding molecules with the general chemical formula of  $CF_3(CF_2)_7SO_2-R$ . The screening health assessment of PFOS, its salts and its precursors containing the  $C_8F_{17}SO_2$  or  $C_8F_{17}SO_3$  moiety covers some 50 substances, many of which are on the Domestic Substances List (see Appendix 1).

The results of a survey of Canadian industry conducted in 2000 under Section 71 of CEPA 1999 to determine the manufacture, import, export and uses of specific PFAs, their derivatives and polymers indicated that there was no known manufacture of PFAs, including PFOS, in Canada. Almost 600 000 kg of PFAs were imported into Canada between 1997 and 2000, PFOS representing only a very small proportion of this total (Environment Canada, 2001). PFOS and its precursors were imported into Canada as chemicals or in various products. As noted elsewhere (OECD, 2002), the principal applications for PFOS and its precursors are for water, oil, soil and grease repellents for use on surface and paper-based applications, such as rugs and carpets, fabric and upholstery and food packaging, as well as use in specialized chemical applications, such as fire-fighting foams, hydraulic fluids, carpet spot removers, mining and oil well surfactants and other specialized chemical formulations. Owing to these use patterns, the exposure of humans to such substances would likely result from contact with, and/or the use of, certain consumer products (3M, 1999a).

The screening health assessment on PFOS, its salts and its precursors containing the  $C_8F_{17}SO_2$  or  $C_8F_{17}SO_3$  moiety is based upon a comparison of the margin between the levels of PFOS in the blood and liver of laboratory animals<sup>1</sup> that are associated with the development of toxicological effects and the levels in the blood and liver of humans. The following considerations were taken into account in developing the approach for the screening health assessment of PFOS, its salts and precursors listed herein:

---

<sup>1</sup> Available data from identified epidemiological studies were considered inadequate for such analysis.

- Chemical, environmental and metabolic processes can lead to the removal of the perfluorinated moiety, ultimately yielding PFOS (3M, 1999a,b).
- PFOS is persistent and is not further degraded or metabolically converted to other compounds (3M, 1999a,b).
- On the basis of CATABOL modelling (Mekenyan and Dimitrov, 2002), the substances listed in Appendix 1 of this assessment were considered to have the potential to biodegrade to PFOS.
- Since PFOS is likely the ultimate perfluorinated degradation or metabolic product of the group of substances listed in Appendix 1, the level of this compound in human tissue provides a useful indicator of exposure to this group of substances from all potential sources.
- Biomonitoring in humans (and animal species) has focused on PFOS, which has been detected in the blood of non-occupationally exposed humans in North America and Europe.
- The toxicity profile of those PFOS precursors examined here (see table below) appears to be generally similar to that of PFOS itself. Available data indicate that effects associated with the PFOS precursors occur at exposures that are similar to or slightly higher than those for PFOS.

Toxicological data relevant to this screening health assessment report were identified for the following substances on the Domestic Substances List (see following table), and these data were used as the basis for assessing the PFOS precursors listed in Appendix 1:

Substance	Designation	CAS No.
1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluoro-(perfluorooctane sulfonate)	PFOS	2795-39-3 (potassium salt) 29081-56-9 (ammonium salt) 70225-14-8 (diethanolamine salt)
1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluoro-N-(2-hydroxyethyl)-(N-ethylperfluorooctane sulfonamidoethanol)	N-EtFOSE	1691-99-2
1-Octanesulfonamide, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluoro-N-(2-hydroxyethyl)-N-methyl-(N-methylperfluorooctane sulfonamidoethanol)	N-MeFOSE	24448-09-7
1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluoro-(N-ethyl perfluorooctane sulfonamide)	N-EtFOSA	4151-50-2
1-Octanesulfonamide, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluoro-N-methyl-(N-methyl perfluorooctane sulfonamide)	N-MeFOSA	31506-32-8
Glycine, N-ethyl-N-[(heptafluorooctyl)-sulfonyl]-, potassium salt (potassium-N-ethyl-N((heptafluorooctyl)-sulfonyl)-glycinate)	PFOSAA	2991-51-7
Ethanaminium, N,N,N-trimethyl-2-[(2-methyl-1-oxo-2-propenyl)oxy]-, chloride, polymer with 2-ethoxyethyl 2-propenoate, 2-[[[(heptafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate and oxiranylmethyl 2-methyl-2-propenoate		92265-81-1

## Exposure Assessment and Hazard Characterization

Although the identified data for some of these substances (see table above) were variable and limited, the available information indicates that toxicological effects of these precursors of PFOS are similar to those of PFOS itself (Table 1). Moreover, based upon the data identified, health-related effects associated with exposure to these substances would appear to be somewhat less severe and/or are observed at higher exposures (doses) than those associated with exposure to PFOS itself.

Studies considered critical to the screening health assessment of PFOS, its salts and its precursors (i.e., those with lowest effect levels) are long-term repeated-dose investigations conducted with rodents and primates. In rats receiving PFOS in the diet for 2 years, histopathological effects in the liver were observed in males and females at intakes as low as 0.06–0.23 mg/kg-bw per day and 0.07–0.21 mg/kg-bw per day, respectively (Covance Laboratories, Inc., 2002a). The mean levels of PFOS in the serum and liver of males and females with these intakes were 7.6 and 20.2 µg/ml and 26.4 and 55.1 µg/g, respectively, after 2 years of exposure (3M Environmental Laboratory, 2001). Evidence of increased thymic atrophy (in females) and reduced serum high-density lipoprotein, cholesterol, bilirubin and triiodothyronine levels (in males) were observed in monkeys administered 0.03 mg PFOS/kg-bw per day for 26 weeks (Covance Laboratories, Inc., 2002b).<sup>2</sup> The mean levels of PFOS in the serum of the males and females after 26 weeks on study were 15.8 and 13.2 µg/ml, respectively (3M Environmental Laboratory, 2000). The mean levels of PFOS in the liver of males and females after 27 weeks on study were 17.3 and 22.2 µg/g, respectively.

Based upon data from chronic exposure studies, there is evidence for the carcinogenicity of PFOS (increased incidence of hepatocellular adenoma in males and females) and N-EtFOSE (increased incidence of hepatocellular adenoma in females; thyroid follicular cell adenoma in males) in rats. In both of these studies, statistically significant increases in tumour incidence were observed only at the highest doses tested — that is, at doses that were above those associated with the development of non-neoplastic effects in these animals. PFOS and its related substances evaluated here are not genotoxic based upon the results of a wide range of identified *in vitro* and *in vivo* assays.

Identified data included investigations on health effects in workers occupationally exposed to PFOS. Although a significantly elevated risk of bladder cancer has been observed for one group of workers exposed to PFOS, the identified epidemiological studies of workers occupationally exposed to PFOS are considered inadequate to assess the potential of this substance (and its precursors) to induce cancer in humans. Workers were exposed to other substances, and the observed number of cause-specific and overall deaths was relatively small.

---

<sup>2</sup> Also reported in Seacat et al. (2002).



Moreover, the database is inadequate as a basis to assess aspects of weight of evidence of causality, such as consistency.

In other surveys of occupationally exposed workers in which potential relationships between exposure to PFOS (assessed by monitoring serum PFOS levels in workers, which ranged as high as 10 µg/ml) and effects upon clinical chemistries and haematological and hormonal parameters were assessed, consistent associations have not been observed. The sensitivity of these investigations is limited by low participation rates and employee turnover during study periods at certain facilities.

In a recent study providing preliminary analytical data on the occurrence and distribution of selected organic perfluorinated compounds in the blood of 56 volunteer non-occupationally exposed adult Canadians, PFOS was detected in 100% of the samples (Kubwabo et al., 2002). Measured concentrations of PFOS in the serum ranged from 0.0037 to 0.065 µg/ml; the overall mean and 95th-percentile concentrations were 0.0288 and 0.0631 µg/ml, respectively. These levels are similar to those measured in other contemporary biomonitoring studies conducted in the United States and Europe (OECD, 2002).

In one study of 599 children (aged 2–12 years) in the United States conducted between 1994 and 1995, the geometric mean concentration of PFOS in the serum was 37.5 ppb (i.e., 0.0375 µg/ml); the 95th percentile was 97 ppb (0.097 µg/ml), with a small number of samples (<20) exceeding this value. Individual values ranging widely from 7 to 515 ppb (i.e., 0.0067–0.515 µg/ml) (3M Medical Department, 2002).

In view of the physical/chemical properties of PFOS, the substance is not expected to accumulate in breast milk. Although biomonitoring data in very young children have not been identified, PFOS levels in the sera of fetuses and very young offspring do not exceed those of exposed dams in experimental studies (Argus Research Laboratories, 1999e,f).

In analyses of liver samples collected from 30 cadavers in the United States, levels of PFOS ranged from <0.0045 (limit of quantitation) to 0.057 µg/g (Olsen et al., 2003). The mean and geometric mean concentrations were 0.0188 and 0.0152 µg/g, respectively.

Confidence in the effects assessment for PFOS and N-EtFOSE is high, owing to the available database, which covers a wide range of toxicological endpoints. Although confidence in the effects assessment for the remaining PFOS-related compounds is low, due to the lack of identified data, this is mitigated to some degree by the consideration that these compounds are likely converted to PFOS in environmental and biological media. Confidence in the measure of the “internal dose” of PFOS (i.e., levels of PFOS in serum and liver) to assess exposure of the general population to this group of compounds is high. Although the size of the sampled population of individuals in Canada is relatively small, the mean measured concentration is similar to that reported elsewhere for samples collected from other non-occupationally exposed populations in the United States and Europe. Although there may be somewhat less confidence

in the available data on the levels of PFOS in human liver, owing to the small sample size in the single report, use of this exposure metric takes into account aspects related to both the toxicokinetics and metabolism of such substances.

### Conclusion for Human Health

The two laboratory studies considered critical for the margin of exposure analysis are the chronic toxicity study in rats, with a large number of animals per dose group exposed for nearly their lifetime, and the study in which small groups of monkeys (considered as better surrogates for humans) received PFOS for 26 weeks. The levels of PFOS in the serum and liver of these animals at the critical effect level are less than those associated with the critical effect in the F<sub>0</sub> and F<sub>1</sub> rats in the two-generation reproduction/developmental toxicity study.

Most often in the screening health assessment of Existing Substances under CEPA 1999, margins of exposure are based upon a comparison of the doses (intakes) administered to laboratory animals at which substance-induced effects were observed with the upper-bound estimate of human intake. While such estimates of intake for PFOS are included in the supporting working document, they are less reliable as a basis for development of margins than those based on serum levels. In this case, therefore, comparisons have been based upon information on the levels of PFOS in the serum and liver from animals administered PFOS and data from human biomonitoring studies. This obviates the need to take into account the significant uncertainty associated with a determination of the upper-bound estimate of human intake of PFOS, owing to the limited available data on levels of PFOS and precursors in air, foodstuffs, drinking water and breast milk and resulting from contact with household materials treated with such perfluorinated substances. Moreover, the levels of PFOS in human tissues provide a useful indicator of the combined exposure from all sources.

Comparisons of the levels of PFOS in the serum and liver of animals at the critical effect level with levels in serum and/or liver from human adult and children biomonitoring studies are presented in the following table:

Critical study and effect	PFOS dose metric at critical effect	Metric(s) of human exposure to PFOS	Margin of exposure (critical effect/human exposure)
Microscopic changes in the liver of rats (m + f) receiving PFOS in the diet for 2 years <sup>1</sup>	Serum PFOS level: 13.9 µg/ml <sup>2</sup>	Mean serum PFOS level in adults in Canada <sup>3</sup> : 0.028 µg/ml	496
		95th percentile of human serum PFOS level in adults in Canada <sup>3</sup> : 0.0631 µg/ml	220
		Mean serum PFOS level in children in the United States <sup>4</sup> : 0.0375 µg/ml	371

Critical study and effect	PFOS dose metric at critical effect	Metric(s) of human exposure to PFOS	Margin of exposure (critical effect/human exposure)
		<b>95th percentile of serum PFOS level in children in the United States<sup>4</sup>:</b> 0.097 µg/ml	143
	<b>Liver PFOS level:</b> 40.8 µg/g <sup>5</sup>	<b>Mean<sup>6</sup> liver PFOS level:</b> 0.0188 µg/g	2170 <sup>7</sup>
Thymic atrophy (f), reduced serum high-density lipoprotein (m), cholesterol (m), triiodothyronine (m) and total bilirubin (m) in monkeys administered PFOS for 26 weeks <sup>1</sup>	<b>Serum PFOS level:</b> 14.5 µg/ml <sup>8</sup>	<b>Mean serum PFOS level in adults in Canada<sup>3</sup>:</b> 0.028 µg/ml	518
		<b>95th percentile of human serum PFOS level in adults in Canada<sup>3</sup>:</b> 0.0631 µg/ml	230
		<b>Mean serum PFOS level in children in the United States<sup>4</sup>:</b> 0.0375 µg/ml	387
		<b>95th percentile of serum PFOS level in children in the United States<sup>4</sup>:</b> 0.097 µg/ml	149
	<b>Liver PFOS level:</b> 19.8 µg/g <sup>9</sup>	<b>Mean<sup>10</sup> liver PFOS level:</b> 0.0188 µg/g	1053 <sup>11</sup>

<sup>1</sup> Covance Laboratories, Inc. (2002a).

<sup>2</sup> Average of mean levels in males (7.6 µg/ml) and females (20.2 µg/ml).

<sup>3</sup> Kubwabo et al. (2002).

<sup>4</sup> 3M Medical Department (2002).

<sup>5</sup> Average of mean levels in males 26.4 (µg/g) and females (55.1 µg/g).

<sup>6</sup> Mean level of PFOS in livers from 30 cadavers (Olsen et al., 2003).

<sup>7</sup> Published data on 95th percentile not available. Margin of exposure based upon highest level of PFOS in human liver from this study (0.057 µg/g) is 716.

<sup>8</sup> Average of mean levels in males (15.8 µg/ml) and females (13.2 µg/ml) (week 26).

<sup>9</sup> Average of mean levels in males 17.3 (µg/g) and females (22.2 µg/g) (week 27).

<sup>10</sup> Mean level of PFOS in livers from 30 cadavers (Olsen et al., 2003).

<sup>11</sup> Published data on 95th percentile not available. Margin of exposure based upon highest level of PFOS in human liver from this study (0.057 µg/g) is 347.

These margins are considered adequate to address elements of uncertainty, including intraspecies variation, interspecies variation and biological adversity or severity of the effects considered critical here. These margins will also be protective for the increased incidence of tumours observed in the chronic study of PFOS in rats, since the tumours were observed only at doses of PFOS that were higher than those that induced non-neoplastic effects and since the weight of evidence indicates that PFOS (and its precursors) are not genotoxic. While the margins for blood levels in children are somewhat less (approximately 145 for the 95th-percentile values), more appropriate margins for comparison with the effect level from long-term studies are those for adults (approximately 225 for the 95th-percentile values), since they are exposed for a greater portion of their life span. In addition, the critical lowest-observed-effect levels

selected for development of these margins of exposure are very conservative, being about an order of magnitude less than values in other studies (i.e., for effects observed in reproductive studies with rats). The margins are also based on more relevant metrics of exposure to PFOS than dose in experimental studies and deterministic estimates of daily intake in children and adults and, as a result, account for a significant portion of the uncertainties associated with interspecies and intraspecies differences in pharmacokinetics (usually accounted for by 4-fold and 3.2-fold default uncertainty factors, respectively). The higher margins for values in liver, although based on limited data, take into account even a greater proportion of uncertainty in toxicokinetics. The margins also take into account limitations of the database for human exposure. Use of the 95th percentiles for the serum levels is also more conservative than deterministic estimates of exposure, which are based on mean intakes of environmental media.

#### Additional Information Identified After Cut-Off Date

The concentration of PFOS measured in 10 maternal pools of plasma obtained from populations residing in Northern Canada ranged from about 20 to 60 ng/mL; overall mean was 36.9 ng/mL (Tittlemier et al, 2004). The mean level of PFOS in 13 cord blood pools was 16.7 ng/mL. In a study involving pregnant Japanese women, blood PFOS levels were higher than those measured in the matching cord blood samples (Inoue et al., 2004). In a study conducted in the United States, Olsen et al. (2005) reported no statistically significant increase in the median concentrations of PFOS, between 1989 and 2001. The median level in sample collected in 1989, were statistically higher than in samples obtained in 1974. Kannan et al. (2004) reported on the levels of PFOS obtained from samples collected in 10 countries; the overall range of mean concentrations (in males or females) was from 1.7 to 73.2 ng/mL.

Table 1: Summary of health effects information for PFOS and related compounds

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...
Acute toxicity: oral	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = 251 mg/kg-bw</b>  (International Research and Development Corporation, 1978a)  [Additional studies: Hazleton Laboratories America, Inc., 1987a; Hazleton Wisconsin, Inc., 1994a; Corning Hazleton, Inc., 1997a]	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = 1467 mg/kg-bw</b>  (International Research and Development Corporation, 1978b)	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = &gt;500 and &lt;5000 mg/kg-bw</b>  (Riker Laboratories, Inc., 1981b)  [Additional study: Riker Laboratories, Inc., 1987]	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = &gt;1000 and &lt;5000 mg/kg-bw</b>  (Riker Laboratories, Inc., 1979)	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = 350 mg/kg-bw</b>  (Hazleton Laboratories America, Inc., 1985a)  [Additional studies: Riker Laboratories, Inc., 1981c; Hazleton Laboratories America, Inc., 1985b]	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = &gt;0.5 and &lt;5 ml/kg-bw</b>  (Biosearch, Inc., 1978c)  [Additional studies: Biosearch, Inc., 1978a; Hazleton Laboratories America, Inc., 1988a]	<b>Oral LD<sub>50</sub> rat (m/f)</b> <b>LD<sub>50</sub> = &gt;5 g/kg-bw</b>  (Hazleton Wisconsin, Inc., 1991a)
Acute toxicity: dermal						<b>Dermal LD<sub>50</sub> rabbit (m/f) 24-hour covered</b> <b>LD<sub>50</sub> = &gt;2000 mg/kg-bw</b>  (Hazleton Laboratories America, Inc., 1988b)	
Acute toxicity: inhalation	<b>Inhalation LC<sub>50</sub> rat (m/f)</b> <b>LC<sub>50</sub> = 5200 mg/m<sup>3</sup></b>  (Bio/Dynamics, Inc., 1979a)	<b>Inhalation LC<sub>50</sub> rat (m/f)</b> <b>LC<sub>50</sub> = &gt;6.5 g/m<sup>3</sup></b>  (Hazleton Laboratories America, Inc., 1981)				<b>Inhalation LC<sub>50</sub> rat (m/f)</b> <b>LC<sub>50</sub> = &gt;22 g/m<sup>3</sup> and &lt;66 g/m<sup>3</sup></b>  (Bio/Dynamics, Inc., 1979b)	

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...
Irritation: ocular	<p><b>Severe irritation:</b> rabbit 0.1 ml ocular application, washout after 5 or 30 seconds</p> <p>(Riker Laboratories, Inc., 1981a)</p> <p>[Additional studies: <b>mild to moderate irritation:</b> Warf Institute, Inc., 1974, 1975; Hazleton Laboratories America, Inc., 1987b; Hazleton Wisconsin, Inc., 1994b; Corning Hazleton, Inc., 1997b]</p>		<p><b>Minimal irritation:</b> rabbit (f) 0.1 g ocular application</p> <p>(Riker Laboratories, Inc., 1984)</p>	<p><b>No irritation:</b> rabbit 0.1 g ocular application</p> <p>(Biosearch, Inc., 1978b)</p>	<p><b>Minimal irritation:</b> rabbit (f) 0.09 g ocular application</p> <p>(Hazleton Laboratories America, Inc., 1985c)</p> <p>[Additional study: Hazleton Laboratories America, Inc., 1985d]</p>	<p><b>Mild irritation:</b> rabbit 0.1 ml ocular application (unwashed)</p> <p>(Hazleton Laboratories America, Inc., 1988c)</p> <p>[Additional study: Biosearch, Inc., 1978d]</p>	<p><b>Moderate irritation:</b> rabbit 0.1 ml ocular application (unwashed)</p> <p>(Hazleton Wisconsin, Inc., 1991b)</p>
Short-term repeated-dose toxicity	<p><b>Oral gavage LOEL</b> rat (m/f), 28 days <b>LOEL = 3 mg/kg-bw per day</b></p> <p>hepatocellular hypertrophy (m/f); increased relative liver weight (m/f); increased relative kidney weight (f); reduced body weight (f)</p> <p>(NOTOX, 1999)</p> <p>[Additional study: Austin et al., 2003]</p>			<p><b>Dietary LOEL</b> rat (m/f), at least 4 weeks <b>LOEL (m/f) = 2.4–4.1 mg/kg-bw per day</b></p> <p>increased relative liver weight (f), hepatocellular hypertrophy (m) (investigators give LOEL of 35–63 mg/kg-bw per day, ignoring liver effects at lower doses)</p> <p>(Covance Laboratories, Inc., 2000a)</p>			

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...
Subchronic toxicity	<p><b>Oral gavage LOEL</b> rhesus monkey (m/f), 90 days  <b>LOEL = 0.5 mg/kg-bw per day</b></p> <p>clinical signs of toxicity and increased leukocytes</p> <p>(International Research and Development Corporation, 1978e)</p> <p>[Additional study: International Research and Development Corporation, 1978c]</p>	<p><b>Oral diet LOEL</b> rat (m/f), 90 days  <b>LOEL (m) = 2 mg/kg-bw per day</b></p> <p>slight hepatocellular vacuolization; decreased hemoglobin and hematocrit</p> <p>(International Research and Development Corporation, 1978d)</p> <p>[Additional study: International Research and Development Corporation, 1979]</p>		<p><b>Oral diet LOEL</b> rat (m/f), 13 weeks  <b>LOEL (m) = 2 mg/kg-bw per day</b></p> <p>increased relative brain, kidney, liver and testis weights; histopathological changes in the liver, reductions in serum cholesterol and triglycerides</p> <p>(Covance Laboratories, Inc., 1999d)</p>			

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...	
Carcinogenicity/ chronic	<p><b>Oral diet LOAEL</b> rat (m/f), 104 weeks <b>LOAEL = 0.06–0.23 mg/kg-bw per day</b></p> <p>increased incidence of non-neoplastic changes in the liver (statistically significant increased incidence of hepatocellular adenoma in males and females at intakes of 0.64–2.21 mg/kg-bw per day)</p> <p>(Covance Laboratories, Inc., 2002a)</p> <p><b>Oral LOEL</b> cynomolgus monkey (m/f), 26 weeks <b>LOEL (m/f) = 0.03 mg/kg-bw per day</b></p> <p><i>m</i>: reduced high-density lipoprotein and triiodothyronine levels, thymic atrophy <i>f</i>: thymic atrophy</p> <p>(Covance Laboratories, Inc., 2002b)</p>	<p><b>Oral diet LOEL</b> rat (m/f) 104-week cancer bioassay with N-EtFOSE narrow range (98.1%) <b>LOEL (m) = 0.86–2.618 mg/kg-bw per day</b> <b>LOEL (f) = 4.213–10.166 mg/kg-bw per day</b></p> <p><i>m/f</i>: increased incidence of histopathological effects in liver <i>f</i>: significant (p &lt; 0.05) reduced serum triglycerides after 104 weeks (statistically significant increased incidences of thyroid follicular adenoma in males at intakes of 3.1–8.72 mg/kg-bw per day and of hepatocellular adenoma in females at intakes of 4.213–10.166 mg/kg-bw per day)</p> <p>(Covance Laboratories, Inc., 2001)</p> <p>[Additional study: Riker Laboratories, Inc., 1983]</p>						
Genotoxicity and related endpoints: <i>in vivo</i>	<p><b>Negative:</b> mouse (m/f) bone marrow micronucleus 950 mg/kg-bw; acute oral gavage</p> <p>(Corning Hazleton, Inc., 1996b)</p>	<p><b>Negative:</b> mouse (m/f) bone marrow micronucleus 2200 mg/kg-bw; acute oral gavage</p> <p>(Corning Hazleton, Inc., 1996a)</p> <p>[Additional studies: Corning Hazleton, Inc., 1993; Hazleton Washington, Inc., 1993a]</p>	<p><b>Negative:</b> mouse (m/f) bone marrow micronucleus, 4000 mg/kg-bw, acute oral gavage</p> <p>(Corning Hazleton, Inc., 1996c)</p>	<p><b>Negative:</b> rat (m/f) bone marrow micronucleus, 5000 mg/kg-bw, acute oral gavage, and rat hepatic unscheduled DNA synthesis <i>in vivo/in vitro</i></p> <p>(Hazleton Washington, Inc., 1993b,c)</p>				
Genotoxicity and related endpoints:	<p><b>Negative:</b> with/without metabolic activation: Ames</p>	<p><b>Negative (-S9), questionable (+S9):</b> mouse</p>	<p><b>Negative:</b> Ames <i>Salmonella</i></p>	<p><b>Negative:</b> with/without</p>	<p><b>Negative:</b> with/without metabolic activation: Ames</p>	<p><b>Negative:</b> with/without</p>		



Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...	
<i>in vitro</i>	<p><i>Salmonella/E. coli</i> mutagenicity, <i>S. cerevisiae</i> mitotic recombinogenicity, rat hepatocyte unscheduled DNA synthesis and human lymphocyte chromosomal aberration <i>in vitro</i> assays</p> <p>(Litton Bionetics, Inc., 1978; SRI International, 1978, 1980, 1981; Covance Laboratories, Inc., 1999a,b,c)</p>	<p>lymphoma L5178Y; <i>in vitro</i> mutation -S9/+S9</p> <p>(NOTOX, 1998)</p> <p>[Additional <b>negative</b> <i>in vitro</i> study: Covance Laboratories, Inc., 2000b]</p>	<p>mutagenicity and Chinese hamster ovary sister chromatid exchange <i>in vitro</i> assays</p> <p>(U.S. EPA, 1989)</p>	<p>metabolic activation: Ames <i>Salmonella</i> mutagenicity, mouse L5178Y lymphoma mutation and human lymphocyte chromosomal aberration <i>in vitro</i> assays</p> <p>(NOTOX, 1994a,b,c)</p>	<p><i>Salmonella</i> mutagenicity and yeast recombination <i>in vitro</i> assays</p> <p>(SRI International, 1985)</p>	<p>metabolic activation: Ames <i>Salmonella</i> mutagenicity and yeast recombination <i>in vitro</i> assays</p> <p>(SRI International, 1982)</p>		
Reproductive/developmental toxicity, rat	<p><b>LOEL maternal/LOEL fetal</b> rat (f) oral gavage, days 6–15 of gestation  <b>LOEL maternal = 5 mg/kg-bw per day</b>  <b>LOEL fetal = 1 mg/kg-bw per day</b></p> <p><i>maternal</i>: decreased weight gain; decreased body weight minus gravid uterine weight; clinical effects  <i>fetal</i>: incomplete skull closure twice that of controls</p> <p>(Hazleton Laboratories America, Inc., 1983b)</p> <p>[Additional studies: Riker Laboratories, Inc., 1980; Argus Research Laboratories, Inc., 1999e,f]</p>	<p><b>LOEL maternal/LOEL fetal</b> rat (f) oral gavage, days 6–17 of gestation  <b>LOEL maternal = 10 mg/kg-bw per day</b>  <b>LOEL fetal = 10 mg/kg-bw per day</b></p> <p><i>maternal</i>: reduced body weight gain  <i>fetal</i>: reduced live fetal body weight and increased skeletal alterations, ossification alterations</p> <p>(Argus Research Laboratories, Inc., 1998)</p> <p>[Additional studies: Riker Laboratories, Inc., 1981d; Hazleton Laboratories America, Inc., 1983a, 1984]</p>						

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...
Reproductive/ developmental toxicity, rat two- generation	<p><b>LOEL F<sub>0</sub>/LOEL F<sub>1</sub>/LOEL F<sub>2</sub>:</b> rat (m/f) oral gavage, F<sub>0</sub> males: from 6 weeks before to the end of mating, F<sub>0</sub> females: from 6 weeks before mating through to the 21st day of lactation (DL 21), F<sub>1</sub> males: from 22 days after birth to the end of mating (started 90 days after birth), F<sub>1</sub> females: from 22 days after birth to DL 21 (for F<sub>2</sub>)</p> <p><b>LOEL F<sub>0</sub> (m) = 0.4 mg/kg-bw per day</b>  <b>LOEL F<sub>0</sub> (f) = 1.6 mg/kg-bw per day</b>  <b>LOEL F<sub>1</sub> (m/f) = 1.6 mg/kg-bw per day</b>  <b>LOEL F<sub>2</sub> (m/f) = &gt;0.4 mg/kg-bw per day</b></p> <p><i>F<sub>0</sub> (m)</i> reduced body weight gains, <i>F<sub>0</sub> (f)</i> reduced body weight gains during precohabitation, <i>F<sub>1</sub> (m/f)</i> significantly reduced litter sizes and both viability and lactation indices; reductions in development, including delayed eye opening, surface righting, pinna unfolding and air righting reflex</p> <p>(Argus Research Laboratories, Inc., 1999a)</p> <p>[Additional study: Argus Research Laboratories, Inc., 2000]</p>	<p><b>LOEL F<sub>0</sub>/LOEL F<sub>1</sub>/LOEL F<sub>2</sub>:</b> rat (m/f) oral gavage, F<sub>0</sub> males: from 28 days before to end of mating, females: from 28 days before through to the 21st day of lactation (DL 21), F<sub>1</sub> males from 22 days after birth to end of mating (started 90 days after birth), F<sub>1</sub> females: from 22 days after birth through to DL 21 (for F<sub>2</sub>)</p> <p><b>LOEL F<sub>0</sub> (m/f) = 5 mg/kg-bw per day</b>  <b>LOEL F<sub>1</sub> (m/f) = 1 mg/kg-bw per day</b>  <b>LOEL F<sub>2</sub> (m/f) = 5 mg/kg-bw per day</b></p> <p><i>F<sub>0</sub></i> reduced body weight gains (m/f); increased relative left testis weight; reduced duration of gestation, <i>F<sub>1</sub></i> reduced body weight gains (m/f), <i>F<sub>2</sub></i> reduced viability and lactation indices; reduced mean litter weight</p> <p>(Argus Research Laboratories, Inc., 1999b)</p>					

Endpoint	PFOS	N-EtFOSE	N-EtFOSA	N-MeFOSE	N-MeFOSA	PFOSAA	Ethanaminium...
Reproductive/ developmental toxicity, rabbit	<p><b>LOEL maternal/ LOEL fetal</b> rabbit (f) oral gavage, days 7–20 of gestation  <b>LOEL maternal = 1.0 mg/kg-bw per day</b>  <b>LOEL fetal = 2.5 mg/kg-bw per day</b></p> <p><i>maternal</i>: reduced body weight gain over entire exposure period  <i>fetal</i>: decreased ossification of sternal centres per fetus per litter; reduced body weight</p> <p>(Argus Research Laboratories, Inc., 1999d)</p>	<p><b>LOEL maternal/LOEL fetal</b> rabbit (f) oral gavage, days 7–20 of gestation  <b>LOEL maternal = 2.5 mg/kg-bw per day</b>  <b>LOEL fetal = &gt;3.75 mg/kg-bw per day</b></p> <p><i>maternal</i>: reduced body weight gain; increased late resorptions and abortions</p> <p>(Argus Research Laboratories, Inc., 1999c)</p> <p>[Additional study: Riker Laboratories, Inc., 1981e]</p>	<p><b>LOEL offspring</b> rabbit (f) oral gavage; days 19–28 of gestation  <b>LOEL offspring = 0.3 mg/kg-bw per day</b></p> <p>increased neonatal mortality throughout pre-weaning period</p> <p>(Stump et al., 1997)</p>				

LOEL = lowest-observed-effect level; LOAEL = lowest-observed-adverse-effect level; LC<sub>50</sub> = median lethal concentration; LD<sub>50</sub> = median lethal dose; m = male; f = female; bw = body weight.

## References

3M. 1999a. Perfluorooctane sulfonate: Current summary of human sera, health and toxicity data. 3M Company, January 21, 1999.

3M. 1999b. The science of organic fluorochemistry. 3M Company, February 5, 1999.

3M Environmental Laboratory. 2000. Determination of the presence and concentration of perfluorooctanesulfonate (PFOS) in liver and serum samples. # FACT TOX-030.

3M Environmental Laboratory. 2001. Determination of the presence and concentration of perfluorooctanesulfonate (PFOS) in liver and serum specimens of CrI:CD<sup>®</sup>(SD) IGS BR rats exposed to perfluorooctane sulfonic acid potassium salt (PFOS T6295). # FACT TOX-002.

3M Medical Department. 2002. Identification of fluorochemicals in human sera. III. Pediatric participants in a group A streptococci clinical trial investigation. Medical Department, 3M Company. # EPI-0011.

Argus Research Laboratories, Inc. 1998. Oral (gavage) developmental toxicity study of N-EtFOSE in rats. # 418-011.

Argus Research Laboratories, Inc. 1999a. Combined oral (gavage) fertility, developmental and perinatal/postnatal reproduction toxicity study of PFOS in rats. # 418-008.

Argus Research Laboratories, Inc. 1999b. Combined oral (gavage) fertility, developmental and perinatal/postnatal reproduction toxicity study of N-EtFOSE in rats. # 418-009.

Argus Research Laboratories, Inc. 1999c. Oral (stomach tube) developmental toxicity study of N-EtFOSE in rabbits. # 418-010.

Argus Research Laboratories, Inc. 1999d. Final report: Oral (stomach tube) developmental toxicity study of PFOS in rabbits. # 418-012.

Argus Research Laboratories, Inc. 1999e. Oral (gavage) pharmacokinetic study of PFOS in rats. # 418-013.

Argus Research Laboratories, Inc. 1999f. Oral (gavage) pharmacokinetic study of PFOS in rats. # 418-015.

Argus Research Laboratories, Inc. 2000. Oral (gavage) cross-fostering study of PFOS in rats. # 418-014.

Austin, M.E., Kasturi, B.S., Barber, M., Kannan, K., MohanKumar, P.S. and MohanKumar, S.M.J. 2003. Neuroendocrine effects of perfluorooctane sulfonate in rats. *Environ. Health Perspect.* 111: 1485–1489.

Bio/Dynamics, Inc. 1979a. An acute inhalation toxicity study of T-2306 CoC in the rat. # 78-7185.

Bio/Dynamics, Inc. 1979b. An acute inhalation toxicity study of T-2307 CoC in the rat. # 78-7186.

Biosearch, Inc. 1978a. Acute oral toxicity — rats. # 77-1108A.

Biosearch, Inc. 1978b. Primary eye irritation study — rabbits. # 78-1161A.

Biosearch, Inc. 1978c. Acute oral toxicity — rats. # 77-1127A.

Biosearch, Inc. 1978d. Primary eye irritation study — rabbits. # 77-1127A.

Corning Hazleton, Inc. 1993. Mutagenicity test on T-5710 in an *in vivo* rat micronucleus assay. # 15516-0-454.

Corning Hazleton, Inc. 1996a. Mutagenicity test on T-6292 in an *in vivo* mouse micronucleus test. # 17384-0-455.

Corning Hazleton, Inc. 1996b. Mutagenicity test on T-6295 in an *in vivo* mouse micronucleus assay. # 17403-0-455.

Corning Hazleton, Inc. 1996c. Mutagenicity test on T-6294 in an *in vivo* mouse micronucleus assay. # 17385-0-455.

Corning Hazleton, Inc. 1997a. Acute oral toxicity study of T-6684 in rats. # CHW61101149.

Corning Hazleton, Inc. 1997b. Final report: Primary eye irritation/corrosion study of T-6684 in rabbits (OECD Guidelines). # 61101151.

Covance Laboratories, Inc. 1999a. *Salmonella–Escherichia coli*/mammalian-microsome reverse mutation assay with PFOS. # 20784-0-409.

Covance Laboratories, Inc. 1999b. Final report: Unscheduled DNA synthesis in rat liver primary cell cultures with PFOS. # 20784-0-447.

Covance Laboratories, Inc. 1999c. Chromosomal aberrations in human whole blood lymphocytes with PFOS. # 20784-0-449.

Covance Laboratories, Inc. 1999d. 13-week dietary toxicity study with T-6314 in rats. # 6329-225.

Covance Laboratories, Inc. 2000a. 4-week range-finding dietary toxicity study with N-methyl perfluorooctanesulfonamido ethanol (N-MeFOSE, T-6314) in rats. # 6329-224.

Covance Laboratories, Inc. 2000b. Final report. L5178Y TK<sup>+</sup> mouse lymphoma forward mutation assay with a confirmatory assay with N-EtFOSE, T-6316. # 20785-0-431 ICH.

Covance Laboratories, Inc. 2001. Final report of the 104 week dietary carcinogenicity study with narrow range (98.1%) N-ethyl perfluorooctanesulfonamido-ethanol (N-EtFOSE) in rats. # 6329-212.

Covance Laboratories, Inc. 2002a. Final report: 104-week dietary chronic toxicity and carcinogenicity study with perfluorooctane sulfonic acid potassium salt (PFOS; T-6295) in rats. # 6329-183.

Covance Laboratories, Inc. 2002b. 26-week capsule toxicity study with perfluorooctane sulfonic acid potassium salt (PFOS T-6295) in cynomolgus monkeys. # 6329-223.

Environment Canada. 2001. Primary report on PFAs from Section 71 survey. Gatineau, Quebec.

Environment Canada. 2003. Technical working document for the environmental screening assessment of perfluorooctane sulfonate (PFOS) and its precursors. Existing Substances Branch, Gatineau, Quebec.

Hazleton Laboratories America, Inc. 1981. Corrected: Final report on acute inhalation toxicity study in rats. # 154-157.

Hazleton Laboratories America, Inc. 1983a. Pilot rat teratology study T-3352. Final report. # 154-159.

Hazleton Laboratories America, Inc. 1983b. Rat teratology study T-3351. # 154-160.

Hazleton Laboratories America, Inc. 1984. Rat teratology study T-3352. # 154-161.

Hazleton Laboratories America, Inc. 1985a. Final report: Acute oral toxicity: method, summary, pathology. # 50202473, sample T-3727.

Hazleton Laboratories America, Inc. 1985b. Final report: Acute oral toxicity: method, summary, pathology. # 50503499, sample T-3752.

Hazleton Laboratories America, Inc. 1985c. Final report, primary eye irritation — method, summary. # 50202473, sample T-3727.

Hazleton Laboratories America, Inc. 1985d. Final report primary eye irritation — method, summary. # 50503501, sample T-3752.

Hazleton Laboratories America, Inc. 1987a. Acute oral toxicity study in rats — method, summary, pathology, raw data appendix. # 70100353, sample T-4016.

Hazleton Laboratories America, Inc. 1987b. Primary eye irritation study in rabbits — method, summary, raw data appendix. # 70100355, sample T-4016.

Hazleton Laboratories America, Inc. 1988a. Acute oral toxicity study in rats (OECD Guidelines). # 70905392.

Hazleton Laboratories America, Inc. 1988b. Acute dermal toxicity study in rabbits (OECD Guidelines). # 70905393.

Hazleton Laboratories America, Inc. 1988c. Primary eye irritation/corrosion study in rabbits (OECD Guidelines). # 70905395.

Hazleton Washington, Inc. 1993a. Genotoxicity test on T-5710 in the *in-vivo/in-vitro* unscheduled DNA synthesis and cell proliferation in rat liver cells. # 15516-0-494.

Hazleton Washington, Inc. 1993b. Mutagenicity test on T-5711 in an *in vivo* rat micronucleus assay. # 15515-0-454.

Hazleton Washington, Inc. 1993c. Genotoxicity test on T-5711.1 (N-MeFOSE) in the *in-vivo/in-vitro* unscheduled DNA synthesis and cell proliferation in rat liver cells. # 15515-0-494.

Hazleton Wisconsin, Inc. 1991a. Acute oral toxicity study of T-5298 in rats (OECD Guidelines). # 01104196.

Hazleton Wisconsin, Inc. 1991b. Primary eye irritation/corrosion study of T-5298 in rabbits (OECD Guidelines). # 01104198.

Hazleton Wisconsin, Inc. 1994a. Acute oral toxicity study of PFOS T-5898 in rats. # 40200468.

Hazleton Wisconsin, Inc. 1994b. Final report: Primary eye irritation/corrosion study of PFOS (T-5898) in rabbits (OECD Guidelines). # 40200470.

Inoue, K., Okada, F., Ito, R., Kato, S., Sasaki, S., Nakajima, S., Uno, A., Saijo, Y., Sata, F., Yoshimura, Y., Kishi, R., and Nakazawa, H. 2004. Perfluorooctane sulfonate (PFOS) and related

perfluorinated compounds in human maternal and cord blood samples: assessment of PFOS exposure in a susceptible population during pregnancy. *Environ. Health Perspec.*, 112: 1204-1207.

International Research and Development Corporation. 1978a. Fluorochemical FC-95, acute oral toxicity (LD<sub>50</sub>) study in rats. # 137-083.

International Research and Development Corporation. 1978b. FM-3422: Acute oral toxicity (LD<sub>50</sub>) study in rats. # 137-084.

International Research and Development Corporation. 1978c. Fluorochemical FC-95, ninety day subacute rat toxicity study. # 137-085.

International Research and Development Corporation. 1978d. Ninety day subacute rat toxicity study, FM-3422. # 137-086.

International Research and Development Corporation. 1978e. Fluorochemical FC-95, ninety day subacute rhesus monkey toxicity study. # 137-092.

International Research and Development Corporation. 1979. Ninety day subacute study in rhesus monkeys. # 137-088.

Kannan, K., Corsolini, S., Falandysz, J., Fillmann, G., Kumar, K.S., Loganathan, B.G., Mohd, M.A., Olivero, J., Van Wouwe, N., Yang, J.H., and Aldoust, K.M. 2004. Perfluorooctanesulfonate and related fluorochemicals in human blood from several countries. *Environ. Sci. Technol.*, 38:4489-95

Kubwabo, C., Vais, N. and Benoit, F. 2002. A pilot study on the determination of the level of perfluorooctanesulfonate and other organic fluorochemicals in human blood. Chemistry Research Division, Safe Environments Programme, Health Canada (unpublished).

Litton Bionetics, Inc. 1978. Mutagenicity evaluation of T-2014 CoC in the Ames *Salmonella*/microsome plate test. # 20838.

Mekenyan, O. and Dimitrov, S. 2002. PFOS metabolic pathways and metabolic distributions: Generated by catabolic simulator (2001–2002). Results compiled and edited by P. Robinson, Environment Canada, June 2002 [cited in Environment Canada, 2003].

NOTOX. 1994a. Evaluation of the mutagenic activity of T-5874 in the Ames *Salmonella*/microsome test (with independent repeat). # 115932.

NOTOX. 1994b. Evaluation of the mutagenic activity of T-5874 in an *in vitro* mammalian cell gene mutation test with L5178Y mouse lymphoma cells (with independent repeat). # 115921.



NOTOX. 1994c. Evaluation of the ability of T-5874 to induce chromosome aberrations in cultured peripheral human lymphocytes (with independent repeat). # 115919.

NOTOX. 1998. Evaluation of the mutagenic activity of T-6906 in an *in vitro* mammalian cell gene mutation test with L5178Y mouse lymphoma cells (with independent repeat). # 223458.

NOTOX. 1999. Exploratory 28-day oral toxicity study with telomer alcohol, telomer acrylate, [redacted confidential business information], PFHS and PFOS (positive control) by daily gavage in the rat followed by a 14/28-day recovery period. # 242933.

OECD (Organisation for Economic Co-operation and Development). 2002. Draft assessment of perfluorooctane sulfonate and its salts. Complete assessment. September 16, 2002.

Olsen, G.W., Hansen, K.J., Stevenson, L.A., Burriss, J.M. and Mandel, J.H. 2003. Human donor liver and serum concentrations of perfluorooctanesulfonate and other perfluorochemicals. *Environ. Sci. Technol.* 37: 888–891.

Olsen, G.W., Huang, H-Y., Helzlsouer, K.J., Hansen, K.J., Butenhoff, J.L. and Mandel, J.H. 2005. Historical comparison of perfluorooctanesulfonate, perfluorooctanoate and other perfluorochemicals in human blood. *Environ. Health Perspec.* 113: 539-545.

Riker Laboratories, Inc. 1979. Acute oral toxicity screen with T2574CoC in albino rats. # 097AR0037.

Riker Laboratories, Inc. 1980. Developmental studies in female rats exposed to FC-95. # 0680TR008.

Riker Laboratories, Inc. 1981a. Acute ocular irritation test with T-2997CoC in albino rabbits. # 0882EB0009.

Riker Laboratories, Inc. 1981b. Acute oral toxicity screen with T-3066CoC in albino rats. # 0981AR0146.

Riker Laboratories, Inc. 1981c. Acute oral toxicity screen with T-3065CoC in albino rats. # 0981AR0145.

Riker Laboratories, Inc. 1981d. Oral teratology study of FM-3422 in rats. # 0680TR0010.

Riker Laboratories, Inc. 1981e. Oral teratology study of T-2999CoC in rabbits. # 0681TB0212.

Riker Laboratories, Inc. 1983. Two year oral (diet) carcinogenicity study of fluorochemical FM-3924 in rats. # 0281CR0012.

Riker Laboratories, Inc. 1984. Acute ocular irritation test with T-3608 in albino rabbits. # 0984EB0367.

Riker Laboratories, Inc. 1987. Acute oral toxicity screen with T-3067CoC in albino rats. # 0981AR0147.

Seacat, A.M., Thomford, P.J., Hansen, K.J., Olsen, G.W., Case, M.T. and Butenhoff, J.L. 2002. Subchronic toxicity studies on perfluorooctanesulfonate potassium salt in cynomolgus monkeys. *Toxicol. Sci.* 68: 249–264.

SRI International. 1978. *In vitro* microbiological mutagenicity assays of 3M Company compounds T-2247 CoC and T-2248 CoC. # LSC-4442-016.

SRI International. 1980. *In vitro* microbiological mutagenicity assays of 3M Company's compound T-2816CoC. # LSC-8958.

SRI International. 1981. *In vitro* microbiological mutagenicity assays of 3M Company's compound T-2997CoC. # LSC-8958

SRI International. 1982. *In vitro* microbiological mutagenicity assays of 3M Company's compound T-3209CoC. # LSC-3145.

SRI International. 1985. *In vitro* microbiological mutagenicity assays of 3M Company's compound T-3727 and T-3752. # LSC-3145.

Stump, D.G., Nemeč, M.D., Holsom, J.F., Piccirillo, V.J. and Mare, J.T. 1997. Study of effects of sulfuramid on pre- and postnatal development, maturation and fertility in the rabbit. *Toxicologist* 36: 357.

Tittlemier, S.A., Ryan, J.J. and van Oostdam, J. 2004. Presence of Anionic Perfluorinated Organic Compounds in Plasma Collected from Northern Canadian Populations. *Organohalogen Compounds*, 64:4009-4014.

U.S. EPA (U.S. Environmental Protection Agency). 1989. Pesticides fact sheet. Office of Pesticides and Toxic Substances. OPTS # 205, 540/FS-89-060.

Warf Institute, Inc. 1974. Primary skin and primary eye irritation assessment of FC-95. #4102871.

Warf Institute, Inc. 1975. Dermal and ocular irritation of PFOS (T-1166) in rabbits. # 5011023.

# **APPENDIX 1**

## **PFOS AND RELATED SUBSTANCES**

CAS No.	Chemical name	Molecular formula
N/A	1-Octanesulfonate, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	$C_8F_{17}SO_3^-$
1691-99-2	1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-N-(2-hydroxyethyl)-	$C_{12}H_{10}F_{17}NO_3S$
2250-98-8	1-Octanesulfonamide, N,N',N''-[phosphinylidynetris(oxy-2,1-ethanediyl)]tris[N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	$C_{36}H_{27}F_{51}N_3O_{10}PS_3$
2795-39-3	1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, potassium salt	$C_8HF_{17}O_3S \cdot K$
2991-51-7	Glycine, N-ethyl-N-[(heptadecafluorooctyl)sulfonyl]-, potassium salt	$C_{12}H_8F_{17}NO_4S \cdot K$
4151-50-2	1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	$C_{10}H_6F_{17}NO_2S$
24448-09-7	1-Octanesulfonamide, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-N-(2-hydroxyethyl)-N-methyl-	$C_{11}H_8F_{17}NO_3S$
29081-56-9	1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, ammonium salt	$C_8HF_{17}O_3S \cdot H_3N$
29117-08-6	Poly(oxy-1,2-ethanediyl), $\alpha$ -[2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl]- $\omega$ -hydroxy-	$(C_2H_4O)_n C_{12}H_{10}F_{17}NO_3S$
30381-98-7	1-Octanesulfonamide, N,N-[phosphinicobis(oxy-2,1-ethanediyl)]bis[N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, ammonium salt	$C_{24}H_{19}F_{34}N_2O_8PS_2 \cdot H_3N$
31506-32-8	1-Octanesulfonamide, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-N-methyl-	$C_9H_4F_{17}NO_2S$
25268-77-3	2-Propenoic acid, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl ester	$C_{14}H_{10}F_{17}NO_4S$
423-82-5	2-Propenoic acid, 2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester	$C_{15}H_{12}F_{17}NO_4S$
38006-74-5	1-Propanaminium, 3-[[heptadecafluorooctyl)sulfonyl]amino]-N,N,N-trimethyl-, chloride	$C_{14}H_{16}F_{17}N_2O_2S \cdot Cl$
52550-45-5	Poly(oxy-1,2-ethanediyl), $\alpha$ -[2-[[heptadecafluorooctyl)sulfonyl]propylamino]ethyl]- $\omega$ -hydroxy-	$(C_2H_4O)_n C_{13}H_{12}F_{17}NO_3S$
56773-42-3	Ethanaminium, N,N,N-triethyl-, salt with 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-1-octanesulfonic acid (1:1)	$C_8H_{20}N \cdot C_8F_{17}O_3S$
57589-85-2	Benzoic acid, 2,3,4,5-tetrachloro-6-[[[3-[[heptadecafluorooctyl)sulfonyl]oxy]phenyl]amino]carbonyl]-, monopotassium salt	$C_{22}H_6Cl_4F_{17}NO_6S \cdot K$
67939-88-2	1-Octanesulfonamide, N-[3-(dimethylamino)propyl]-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, monohydrochloride	$C_{13}H_{13}F_{17}N_2O_2S \cdot ClH$
67969-69-1	1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-N-[2-(phosphonooxy)ethyl]-, diammonium salt	$C_{12}H_{11}F_{17}NO_6PS_2 \cdot H_3N$
68298-11-3	1-Propanaminium, 3-[[heptadecafluorooctyl)sulfonyl](3-sulfopropyl)amino]-N-(2-hydroxyethyl)-N,N-dimethyl-, hydroxide, inner salt	$C_{18}H_{23}F_{17}N_2O_6S_2$
68298-62-4	2-Propenoic acid, 2-[butyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester, telomer with 2-[butyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, methyloxirane polymer with oxirane di-2-propenoate, methyloxirane polymer with oxirane mono-2-propenoate and 1-octanethiol	$(C_{17}H_{16}F_{17}NO_4S \cdot C_{16}H_{16}F_{15}NO_4S \cdot W_{99} \cdot W_{99})_x \cdot C_8H_{18}S$

CAS No.	Chemical name	Molecular formula
68298-78-2	2-Propenoic acid, 2-methyl-, 2-[[[5-[[[2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethoxy]carbonyl]amino]-2-methylphenyl]amino]carbonyl]oxy]propyl ester, telomer with butyl 2-propenoate, 2-[[[5-[[[2-[ethyl[(nonafluorobutyl)sulfonyl]amino]ethoxy]carbonyl]amino]-2-methylphenyl]amino]carbonyl]oxy]propyl 2-methyl-2-propenoate, 2-[[[5-[[[2-[ethyl[(pentadecafluoroheptyl)sulfonyl]amino]ethoxy]carbonyl]amino]-2-methylphenyl]amino]carbonyl]oxy]propyl 2-methyl-2-propenoate, 2-[[[5-[[[2-[ethyl[(tridecafluorohexyl)sulfonyl]amino]ethoxy]carbonyl]amino]-2-methylphenyl]amino]carbonyl]oxy]propyl 2-methyl-2-propenoate, 2-[[[5-[[[2-[ethyl[(undecafluoropentyl)sulfonyl]amino]ethoxy]carbonyl]amino]-2-methylphenyl]amino]carbonyl]oxy]propyl 2-methyl-2-propenoate, 2-[[[5-[[[2-[ethyl[(heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate and 1-octanethiol	$(C_{28}H_{28}F_{17}N_3O_8 \cdot C_{27}H_{28}F_{15}N_3O_8 \cdot C_{26}H_{28}F_{13}N_3O_8 \cdot C_{25}H_{28}F_{11}N_3O_8 \cdot C_{24}H_{28}F_9N_3O_8 \cdot C_{14}H_{10}F_{17}NO_4 \cdot C_{13}H_{10}F_{15}NO_4 \cdot C_{12}H_{10}F_{13}NO_4 \cdot C_{11}H_{10}F_{11}NO_4 \cdot C_{10}H_{10}F_9NO_4 \cdot C_7H_{12}O_2)_x \cdot C_8H_{18}S$
68329-56-6	2-Propenoic acid, eicosyl ester, polymer with 2-[[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate, hexadecyl 2-propenoate, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate and octadecyl 2-propenoate	$(C_{23}H_{44}O_2 \cdot C_{21}H_{40}O_2 \cdot C_{19}H_{36}O_2 \cdot C_{14}H_{10}F_{17}NO_4 \cdot C_{13}H_{10}F_{15}NO_4 \cdot C_{12}H_{10}F_{13}NO_4 \cdot C_{11}H_{10}F_{11}NO_4 \cdot C_{10}H_{10}F_9NO_4)_x$
68555-90-8	2-Propenoic acid, butyl ester, polymer with 2-[[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate and 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate	$(C_{14}H_{10}F_{17}NO_4 \cdot C_{13}H_{10}F_{15}NO_4 \cdot C_{12}H_{10}F_{13}NO_4 \cdot C_{11}H_{10}F_{11}NO_4 \cdot C_{10}H_{10}F_9NO_4 \cdot C_7H_{12}O_2)_x$
68555-91-9	2-Propenoic acid, 2-methyl-, 2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester, polymer with 2-[ethyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate and octadecyl 2-methyl-2-propenoate	$(C_{22}H_{42}O_2 \cdot C_{16}H_{14}F_{17}NO_4 \cdot C_{15}H_{14}F_{15}NO_4 \cdot C_{14}H_{14}F_{13}NO_4 \cdot C_{13}H_{14}F_{11}NO_4 \cdot C_{12}H_{14}F_9NO_4)_x$
68555-92-0	2-Propenoic acid, 2-methyl-, 2-[[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl ester, polymer with 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-methyl-2-	$(C_{22}H_{42}O_2 \cdot C_{15}H_{12}F_{17}NO_4 \cdot C_{14}H_{12}F_{15}NO_4 \cdot C_{13}H_{12}F_{13}NO_4 \cdot C_{12}H_{12}F_{11}NO_4 \cdot C_{11}H_{12}F_9NO_4)_x$

CAS No.	Chemical name	Molecular formula
	propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate and octadecyl 2-methyl-2-propenoate	
68586-14-1	2-Propenoic acid, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl ester, telomer with 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, $\alpha$ -(2-methyl-1-oxo-2-propenyl)- $\omega$ -hydroxypoly(oxy-1,2-ethanediyl), $\alpha$ -(2-methyl-1-oxo-2-propenyl)- $\omega$ -[(2-methyl-1-oxo-2-propenyl)oxy]poly(oxy-1,2-ethanediyl), 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate and 1-octanethiol	$(C_{14}H_{10}F_{17}NO_4S \cdot C_{13}H_{10}F_{15}NO_4S \cdot C_{12}H_{10}F_{13}NO_4S \cdot C_{11}H_{10}F_{11}NO_4S \cdot C_{10}H_{10}F_9NO_4S \cdot (C_2H_4O)_n \cdot C_8H_{10}O_3 \cdot (C_2H_4O)_n \cdot C_4H_6O_2)_x \cdot C_8H_{18}S$
68649-26-3	1-Octanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-N-(2-hydroxyethyl)-, reaction products with N-ethyl-1,1,2,2,3,3,4,4,4-nonafluoro-N-(2-hydroxyethyl)-1-butanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,7-pentadecafluoro-N-(2-hydroxyethyl)-1-heptanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,6-tridecafluoro-N-(2-hydroxyethyl)-1-hexanesulfonamide, N-ethyl-1,1,2,2,3,3,4,4,5,5,5-undecafluoro-N-(2-hydroxyethyl)-1-pentanesulfonamide, polymethylenepolyphenylene isocyanate and stearyl alc.	N/A
68867-62-9	2-Propenoic acid, 2-methyl-, 2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester, telomer with 2-[ethyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 1-octanethiol and $\alpha$ -(1-oxo-2-propenyl)- $\omega$ -methoxypoly(oxy-1,2-ethanediyl)	$(C_{16}H_{14}F_{17}NO_4S \cdot C_{15}H_{14}F_{15}NO_4S \cdot C_{14}H_{14}F_{13}NO_4S \cdot C_{13}H_{14}F_{11}NO_4S \cdot C_{12}H_{14}F_9NO_4S \cdot (C_2H_4O)_n \cdot C_4H_6O_2)_x \cdot C_8H_{18}S$
68877-32-7	2-Propenoic acid, 2-methyl-, 2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester, polymer with 2-[ethyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate, 2-[ethyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-methyl-2-propenoate and 2-methyl-1,3-butadiene	$(C_{16}H_{14}F_{17}NO_4S \cdot C_{15}H_{14}F_{15}NO_4S \cdot C_{14}H_{14}F_{13}NO_4S \cdot C_{13}H_{14}F_{11}NO_4S \cdot C_{12}H_{14}F_9NO_4S \cdot C_5H_8)_x$
68891-96-3	Chromium, diaquatetrachloro[ $\mu$ -[N-ethyl-N-[(heptadecafluorooctyl)sulfonyl]glycinato-O':O"]] $\mu$ -hydroxybis(2-methylpropanol)di-	$C_{18}H_{28}Cl_4Cr_2F_{17}NO_9S$
68958-61-2	Poly(oxy-1,2-ethanediyl), $\alpha$ -[2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl]- $\omega$ -methoxy-	$(C_2H_4O)_n \cdot C_{13}H_{12}F_{17}NO_3S$
70225-14-8	1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, compd. with 2,2-iminobis[ethanol] (1:1)	$C_8HF_{17}O_3S \cdot C_4H_{11}NO_2$
70776-36-2	2-Propenoic acid, 2-methyl-, octadecyl ester, polymer with 1,1-dichloroethene, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate, N-(hydroxymethyl)-2-propenamide, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, 2-	$(C_{22}H_{42}O_2 \cdot C_{14}H_{10}F_{17}NO_4S \cdot C_{13}H_{10}F_{15}NO_4S \cdot C_{12}H_{10}F_{13}NO_4S \cdot C_{11}H_{10}F_{11}NO_4S \cdot C_{10}H_{10}F_9NO_4S \cdot C_4H_7NO_2 \cdot C_2H_2Cl_2)_x$

CAS No.	Chemical name	Molecular formula
	[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate and 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate	
71487-20-2	2-Propenoic acid, 2-methyl-, methyl ester, polymer with ethenylbenzene, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl 2-propenoate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl 2-propenoate and 2-propenoic acid	$(C_{14}H_{10}F_{17}NO_4S \cdot C_{13}H_{10}F_{15}NO_4S \cdot C_{12}H_{10}F_{13}NO_4S \cdot C_{11}H_{10}F_{11}NO_4S \cdot C_{10}H_{10}F_9NO_4S \cdot C_8H_8 \cdot C_5H_8O_2 \cdot C_3H_4O_2)_x$
92265-81-1	Ethanaminium, N,N,N-trimethyl-2-[(2-methyl-1-oxo-2-propenyl)oxy]-, chloride, polymer with 2-ethoxyethyl 2-propenoate, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl 2-propenoate and oxiranylmethyl 2-methyl-2-propenoate	$(C_{14}H_{10}F_{17}NO_4S \cdot C_9H_{18}NO_2 \cdot C_7H_{12}O_3 \cdot C_7H_{10}O_3 \cdot Cl)_x$
94313-84-5	Carbamic acid, [5-[[[2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethoxy]carbonyl]amino]-2-methylphenyl]-, 9-octadecenyl ester, (Z)-	$C_{38}H_{50}F_{17}N_3O_6S$
98999-57-6	Sulfonamides, C <sub>7-8</sub> -alkane, perfluoro, N-methyl-N-[2-[(1-oxo-2-propenyl)oxy]ethyl], polymers with 2-ethoxyethyl acrylate, glycidyl methacrylate and N,N,N-trimethyl-2-[(2-methyl-1-oxo-propenyl)oxy]ethanaminium chloride	$(C_{14}H_{10}F_{17}NO_4S \cdot C_9H_{18}NO_2 \cdot C_7H_{12}O_3 \cdot C_7H_{10}O_3 \cdot Cl)_x$
178094-69-4	1-Octanesulfonamide, N-[3-(dimethyloxidoamino)propyl]-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, potassium salt	$C_{13}H_{12}F_{17}N_2O_3S \cdot K$
N/A	2-(Perfluoro-N-methyl-C <sub>4-8</sub> -1-alkanesulfonamido)ethyl esters of trimers of C <sub>18</sub> unsaturated fatty acids	N/A
29457-72-5	1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-, lithium salt	$C_8HF_{17}O_3S \cdot Li$
68909-15-9	2-Propenoic acid, eicosyl ester, polymers with branched octyl acrylate, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl acrylate, 2-[methyl[(nonafluorobutyl)sulfonyl]amino]ethyl acrylate, 2-[methyl[(pentadecafluoroheptyl)sulfonyl]amino]ethyl acrylate, 2-[methyl[(tridecafluorohexyl)sulfonyl]amino]ethyl acrylate, 2-[methyl[(undecafluoropentyl)sulfonyl]amino]ethyl acrylate, polyethylene glycol acrylate Me ether and stearyl acrylate	$(C_{23}H_{44}O_2 \cdot C_{21}H_{40}O_2 \cdot C_{14}H_{10}F_{17}NO_4S \cdot C_{13}H_{10}F_{15}NO_4S \cdot C_{12}H_{10}F_{13}NO_4S \cdot C_{11}H_{10}F_{11}NO_4S \cdot C_{10}H_{10}F_9NO_4S \cdot (C_2H_4O)_n \cdot C_4H_6O_2 \cdot \text{Unspecified})_x$
148684-79-1	Sulfonamides, C <sub>4-8</sub> -alkane, perfluoro, N-(hydroxyethyl)-N-methyl, reaction products with 1,6-diisocyanatohexane homopolymer and ethylene glycol	N/A
30295-51-3	1-Octanesulfonamide, N-[3-(dimethyloxidoamino)propyl]-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	N/A
91081-99-1	Sulfonamides, C <sub>4-8</sub> -alkane, perfluoro, N-(hydroxyethyl)-N-methyl, reaction products with epichlorohydrin, adipates (esters)	N/A
N/A	Fatty acids, C <sub>18</sub> -unsatd., dimers, 2-[methyl[(perfluoro-C <sub>4-8</sub> -alkyl)sulfonyl]amino]ethyl esters	N/A
68081-83-4	Carbamic acid, (4-methyl-1,3-phenylene)bis-, bis[2-[ethyl[(perfluoro-C <sub>4-8</sub> -alkyl)sulfonyl]amino]ethyl] ester	
68608-14-0	Sulfonamides, C <sub>4-8</sub> -alkane, perfluoro, N-ethyl-N-(hydroxyethyl), reaction products with 1,1'-methylenebis[4-isocyanatobenzene]	

CAS No.	Chemical name	Molecular formula
307-35-7	1-Octanesulfonyl fluoride, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	N/A
376-14-7	2-Propenoic acid, 2-methyl-, 2-[ethyl[(heptadecafluorooctyl)sulfonyl]amino]ethyl ester	N/A
14650-24-9	2-Propenoic acid, 2-methyl-, 2-[[heptadecafluorooctyl)sulfonyl]methylamino]ethyl ester	N/A
94133-90-1	1-Propanesulfonic acid, 3-[[3-(dimethylamino)propyl][(heptadecafluorooctyl)sulfonyl]amino]-2-hydroxy-, monosodium salt	N/A
127133-66-8	2-Propenoic acid, 2-methyl-, polymers with Bu methacrylate, lauryl methacrylate and 2-[methyl[(perfluoro-C <sub>4-8</sub> -alkyl)sulfonyl]amino]ethyl methacrylate	N/A
179005-06-2	Sulfonamides, C <sub>4-8</sub> -alkane, perfluoro, N-[3-(dimethyloxidoamino)propyl], potassium salts	N/A
179005-07-3	Sulfonamides, C <sub>4-8</sub> -alkane, perfluoro, N-[3-(dimethyloxidoamino)propyl]	N/A
ROF	Residual Organic Fluorochemicals (impurities)	N/A
1763-23-1	1-Octanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluoro-	C <sub>8</sub> HF <sub>17</sub> O <sub>3</sub> S

N/A = not available; Me = methyl; Bu = butyl