

Canada’s Provincial and Territorial Routine (and Catch-up) Vaccination Schedule for Infants and Children (as of December 2025)

This table summarizes the current routine vaccination schedule for infants and children in all provinces and territories across Canada. It is updated regularly in collaboration with the Canadian Immunization Committee (CIC)

| VACCINES | | Provincial & Territorial Vaccination Schedules | | | | | | | | | | | | |
|-----------------|---|---|----------------------|---|---|--|--|----------------------|----------------------|---|---|---|--|-----------------------|
| Abbreviations | Description | BC | AB | SK | MB | ON | QC ² | NB | NS | PE | NL | YT | NT | NU |
| DTaP-IPV-Hib | Diphtheria, tetanus, acellular pertussis, inactivated polio virus, Haemophilus influenzae type B vaccine | Age: 18 mos | Age: 18mos | Age: 2,4,6,18mos | Age: 2,4,6,18mos | Age: 2,4,6,18mos | Age: 12mos | Age: 2,4,6,18mos | Age: 2,4,6,18mos | Age: 18mos | Age: 2,4,6,18mos | Age: 18mos | Age: 2,4,6,18mos | Age: 2,4,6,18 mos |
| DTaP-HB-IPV-Hib | Diphtheria, tetanus, acellular pertussis, hepatitis B, inactivated polio virus, Haemophilus influenzae type B vaccine | Age: 2,4,6mos | Age: 2,4,6mos | | | | Age: 2,4mos | | | Age: 2,4,6mos | | Age: 2,4,6mos | | |
| Tdap-IPV | Tetanus, diphtheria (reduced toxoid), acellular pertussis (reduced toxoid), inactivated polio virus vaccine | Age: 4-6 yrs | Age: 4 yrs | Age: 4-6yrs | Age: 4-6 yrs | Age: 4-6 yrs | Age: 4-6 yrs | Age: 4 yrs | Age: 4-6yrs | Age: 4-5 yrs | Age: 4-6yrs ³ | Age: 4-6yrs | Age: 4-6yrs ³ | Age: 4-6 yrs |
| Tdap | Tetanus, diphtheria (reduced toxoid), acellular pertussis (reduced toxoid) vaccine | Grade 9 | Grade 9 | Grade 8 | Grade 8 or 9 | Age: 14-16yrs | | Grade 7 | Grade 7 | Grade 9 | Grades 7 and 8 | Grade 9 | Grade 7 | Grade 6 |
| Td | Tetanus, diphtheria (reduced toxoid) vaccine | | | | | | 3 rd year of high school | | | | | | | |
| BCG | Bacille-Calmette-Guérin vaccine | | | | | | | | | | | | Age :1 mos | Age :1 mos |
| HAHB | Hepatitis A and B vaccine | | | | | | Age: 18 mos | | | | | | | |
| HA | Hepatitis A vaccine | Age: 6, 18 mos, Indigenous children only | | | | | 4 th year of primary school | | | | | | | |
| HB | Hepatitis B vaccine | HB is provided in a 3-dose combination vaccine (DTaP-HB-IPV-Hib) in infancy | Grade 6 (2 doses) | Grade 6 (2 doses) | Grade 6 (2 doses) | Grade 7 (2 doses) | HB is provided in a 2-dose combination vaccine (DTaP-HB-IPV-Hib) in infancy 1-dose of HAHB at 18 mos | Age: At birth,2,6mos | Grade 7 (2 doses) | HB is provided in a 3-dose combination vaccine (DTaP-HB-IPV-Hib) in infancy | Grade 6 (2 doses) | HB is provided in a 3-dose combination vaccine(DTaP-HB-IPV-Hib) in infancy | Age: At birth,1,6 mos | Age: At birth,1,9 mos |
| MMR | Measles, mumps, rubella vaccine | Age: 12 mos | | | | Age: 12mos | | | | | | Age: 12mos | | |
| Var | Varicella vaccine | Age: 12 mos | | | | Age: 15mos | | | | | | Age: 12mos | | Grade 6 (catch-up) |
| MMR-V | Measles, mumps, rubella, and varicella vaccine | Age: 4-6 yrs 2 nd dose | Age: 12, 18 mos | Age: 12,18mos | Age: 12mos, 4-6yrs | Age: 4-6 yrs 2 nd dose | Age: 12,18mos | Age: 12,18mos | Age: 12,18 mos | Age: 12,18mos | Age: 12,18mos | Age: 4-6 years 2 nd dose | Age: 12,18mos | Age: 12, 18mos |
| Men-C-C | Monovalent conjugate meningococcal vaccines | Age: 2,12 mos | Age: 4,12mos | Age: 12mos | | Age: 12mos | Age: 18 mos | Age: 12mos | Age: 12mos | Age: 12mos | Age: 12mos | Age: 2,12mos | Age: 2,12mos | Age: 12 mos |
| Men-C-ACYW-135 | Quadrivalent conjugate meningococcal vaccines | Grade 9 | Grade 9 | Grade 8 | Age:12 mos and Grade 6 | Grade 7 | 3 rd year of high school | Grade 9 | Grade 7 | Grade 9 | Grade 4 | Grade 9 | Grade 12 ¹ | Grade 9 |
| Pneu-C-13 | Pneumococcal conjugate, 13-valent vaccine | Age: 2,4,12 mos | | | | | | | | | | | | Age: 2,4,6,15 mos |
| Pneu-C-15 | Pneumococcal conjugate, 15-valent vaccine | | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,12mos | Age: 2,4,6,18mos | |
| Rota | Rotavirus vaccines ⁴ | Age: 2,4 mos | Age: 2,4mos | Age: 2,4,6mos | Age: 2,4mos | Age: 2,4mos | Age: 2,4mos | Age: 2,4,6 mos | Age: 2,4,6 mos | Age: 2,4,6mos | Age: 2,4mos | Age: 2,4,6mos | Age: 2,4,6mos | Age: 2,4,6 mos |
| HPV | Human papillomavirus vaccines | Grade: 6 (2 doses) | Grade 6 (2 doses) | Grade 6 (2 doses) | Grade 6 (2 doses) | Grade 7 (2 doses) | 4 th year of primary school (1 dose) | Grade 7 (1 dose) | Grade 7 (2 doses) | Grade 6 (2 doses) | Grade 6 (2 doses) | Grade 6 (1 dose) | Grade 4, 5, or 6 (2 doses) | Grade 6 (1 dose) |
| Inf | Influenza vaccines | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Not recommended but offered annually to anyone aged 6 mos or older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older | Age: 6 mos and older |
| RSV | Monoclonal antibodies against the respiratory syncytial virus (not a vaccine) | | | All infants entering or born during a RSV season (from October 1 st to March 31) | All infants entering or born during a RSV season (from October 1 st to March 31) | Infants born on or after April 1 st and less than 8 months of age at the time of immunization | All infants born on or after April 1 st of current year | | | All infants (<8 mos) entering or born during a RSV season | Infants and children at high-risk of severe RSV disease | All infants (<9 mos) entering or born during a RSV season | All infants (0-3 mos) entering or born during a RSV season | All infants at birth |

LEGEND

| Abbreviation/ footnote | Definition |
|---------------------------|--|
| yrs | Years (age) |
| mos | Months (age) |
| 1 | If attending post-secondary school or military recruits who will be living in a congregate setting. |
| 2 | A catch-up program in the 3 rd year of high-school evaluates student's vaccination history. Students will be provided any missed vaccination to protect against the following diseases: diphtheria, pertussis, tetanus, polio, measles, mumps, rubella, varicella, meningococcal serogroup C infections, hepatitis A and B, and human papilloma virus. Temporary catch-up schedule: children in the fourth year of primary school will be vaccinated against hepatitis A. |
| 3 | DTaP-IPV-Hib may be substituted for DTaP-IPV in times of shortage for those who are 4-6yrs of age. |
| 4 | There are two rotavirus vaccines authorized in Canada: Rotarix (2 doses) and RotaTeq (3 doses). |
| | Vaccine is not publicly funded in this province/territory. |