

February 26 to March 4, 2017 (Week 09)


## Overall Summary

- Overall, the decline in influenza activity in Canada has been slow compared to previous seasons. Many parts of Canada are still reporting elevated activity in week 09.
- Widespread or localized influenza activity was reported in 29 regions across 11 provinces.
- In week 09, laboratory detections, influenza-like illness, outbreaks and hospitalizations from participating provinces and territories and sentinel networks decreased from the previous week.
- Influenza B activity in Canada is slowly increasing but remain below what has been observed in previous seasons.
- A(H3N2) continues to be the most common type of influenza affecting Canadians.
- The majority of laboratory detections, hospitalizations and deaths have been among adults aged 65+ years.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

## Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In week 09, two regions (one each in NB and NT) reported no influenza or influenza-like illness activity. Sporadic influenza activity was reported in 22 regions across 10 provinces and territories. Localized activity was reported in 25 regions across nine provinces. Widespread activity was reported in four regions (PE, two in BC and one in QC). For more details on a specific region, click on the map.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 09




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

## Laboratory Confirmed Influenza Detections

In week 09, the number of positive tests (2,072) and the percentage of tests positive for influenza (22%) decreased from the previous week. Peak influenza detections occurred in week 02 at 27%. After a decline from the peak in week 03, detections have remained relatively stable (ranging from 22% to 25% in weeks 03 to 09). Influenza A continues to account for the majority of detections; however, influenza B detections have been steadily increasing for the past few weeks. Influenza B activity is very low compared to the same time period in the previous two seasons. For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada (PHAC) website.


**Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2016-17, Week 09**



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

To date, 29,910 laboratory confirmed influenza detections have been reported, of which 97% have been influenza A. Influenza A(H3N2) is the most common subtype detected. For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

**Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2016-17, Week 09**



To date, detailed information on age and type/subtype has been received for 20,914 laboratory-confirmed influenza cases (Table 1). Among cases with reported age and type/subtype information, adults aged 65+ accounted for almost half of the reported influenza cases. Among cases of influenza A(H3N2), adults aged 65+ represented 46% of cases, followed by adults aged 20-64 (33% of cases). In the previous influenza A(H3N2)-predominant season in 2014-15, adults aged 65+ represented 59% of cases and adults aged 20-64 represented 26% of cases.

**Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting<sup>1</sup>, Canada, 2016-17, Week 09**

Age groups (years)	Week (February 26 to March 4, 2017)					Cumulative (August 28, 2016 to March 4, 2017)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>3</sup>		A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>3</sup>		Total	#
0-4	123	0	11	112	8	2349	10	1028	1311	79	2428	12%
5-19	116	0	27	89	10	2131	9	1127	995	104	2235	11%
20-44	100	0	11	89	10	2654	23	1445	1186	100	2754	13%
45-64	132	0	19	113	14	3523	19	1784	1720	134	3657	17%
65+	497	0	74	423	41	9634	10	4523	5101	206	9840	47%
<b>Total</b>	968	0	142	826	83	20291	71	9907	10313	623	20914	100%
<b>Percentage<sup>2</sup></b>	92%	0%	15%	85%	8%	97%	0%	49%	51%	3%		

<sup>1</sup>Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. Cumulative data include updates to previous weeks.

<sup>2</sup>Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

<sup>3</sup>UnS: unsubtype: The specimen was typed as influenza A, but no result for subtyping was available.

x: Suppressed to prevent residual disclosure

Specimens from NT, YT, and NU are sent to reference laboratories in the provinces


## Syndromic/Influenza-like Illness Surveillance

### Healthcare Professionals Sentinel Syndromic Surveillance

In week 09, 1.8% of visits to healthcare professionals were due to influenza-like illness, down from 2.2% in the previous week.

**Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2016-17**

Number of Sentinels Reporting Week 09: 109



Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.


**Are you a primary healthcare practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel? Please visit our [Influenza Sentinel page](#) for more details.**

## Influenza Outbreak Surveillance

In week 09, 43 laboratory confirmed influenza outbreaks were reported (18 less than the previous week). Among the reported outbreaks: 30 were in long-term care (LTC) facilities, four in hospitals and nine in institutional or community (other) settings. Of the outbreaks with known strains or subtypes, 11 were due to influenza A(H3N2), 15 were due to influenza A(UnS) and one outbreak was due to influenza B. An additional outbreak due to ILI was reported in a school.

To date this season, 902 outbreaks have been reported and the majority (67%) have occurred in LTC facilities. Compared to the same period in the most recent previous A(H3N2) predominant season (2014-15), 1,436 outbreaks were reported, of which 74% occurred in LTC facilities.

**Figure 5 – Number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2016-17, Week 09**



<sup>1</sup>All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

## Provincial/Territorial Influenza Hospitalizations and Deaths

In week 09, 225 influenza-associated hospitalizations were reported by participating provinces and territories, down slightly from 245 reported in the previous week<sup>1</sup>. Influenza A accounted for 90% of hospitalizations. The weekly percentage of hospitalizations due to influenza B has been steadily increasing for the past few weeks. The largest proportion of hospitalizations were among adults aged 65+ (67%). A total of nine intensive care unit (ICU) admissions and 15 deaths were reported in week 09.

To date this season, 4,556 hospitalizations have been reported, of which 98% were due to influenza A. Among cases for which the subtype of influenza A was reported, almost all (2517/2527) were influenza A(H3N2). Adults 65+ accounted for 69% of the hospitalizations. A total of 168 ICU admissions and 206 deaths have been reported. The majority of deaths were reported in adults aged 65+ years.

**Table 2 – Cumulative number of hospitalizations, ICU admissions and deaths by age and influenza type reported by participating provinces and territories, Canada, 2016-17, Week 09**

Age Groups (years)	Cumulative (August 28, 2016 to March 4, 2017)						
	Hospitalizations			ICU Admissions		Deaths	
	Influenza A Total	Influenza B Total	Total [# (%)]	Influenza A and B Total	%	Influenza A and B Total	%
0-4	335	14	349 (7%)	10	6%	<5	x%
5-19	191	17	208 (5%)	12	7%	<5	x%
20-44	244	6	250 (5%)	17	10%	<5	x%
45-64	604	17	621 (14%)	45	27%	29	14%
65+	3076	52	3128 (69%)	84	50%	172	83%
<b>Total</b>	<b>4450</b>	<b>106</b>	<b>4556 (100%)</b>	<b>168</b>	<b>100%</b>	<b>206</b>	<b>100%</b>

x: Suppressed to prevent residual disclosure

<sup>1</sup>Note: Influenza-associated hospitalizations are not reported to PHAC by BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.


## Sentinel Hospital Influenza Surveillance

### Pediatric Influenza Hospitalizations and Deaths


In week 09, 14 laboratory-confirmed influenza-associated pediatric ( $\leq 16$  years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. All but two cases were due to influenza A. The number of weekly hospitalizations reported since week 05 have been below the six year average for the same time period (Figure 7).

To date this season, 421 laboratory-confirmed influenza-associated pediatric hospitalizations were reported by the IMPACT network. Children aged 0-23 months accounted for approximately 41% of hospitalizations. Influenza A accounted for 93% (n=386) of the reported hospitalizations, of which 36% (n=138) were influenza A(H3N2) and the remainder were A(UnS). Additionally, 66 intensive care unit (ICU) admissions have been reported. Children aged 0-23 months accounted for 30% of ICU cases followed by children aged 10-16 (26%). A total of 43 ICU cases reported at least one underlying condition or comorbidity. Less than five deaths have been reported this season.

**Figure 6 – Cumulative numbers of pediatric hospitalizations ( $\leq 16$  years of age) with influenza by age-group reported by the IMPACT network, Canada, 2016-17, Week 09**



**Figure 7 – Number of pediatric hospitalizations ( $\leq 16$  years of age) with influenza reported by the IMPACT network, by week, Canada, 2016-17, Week 09**



The shaded area represents the maximum and minimum number of cases reported by week from seasons 2010-11 to 2015-16


The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

## Adult Influenza Hospitalizations and Deaths


In week 09, 48 laboratory-confirmed influenza-associated adult ( $\geq 20$  years of age) hospitalizations were reported by the Canadian Immunization Research Network (CIRN). All but two cases were due to influenza A and the majority of cases (81%) occurred in adults aged 65+. The number of hospitalizations reported weekly by CIRN have been consistently declining since week 07.

To date this season, 1,100 laboratory-confirmed influenza-associated adult ( $\geq 20$  years of age) hospitalizations have been reported by CIRN. Influenza A accounted for 99% of hospitalizations. Adults aged 65+ accounted for 78% of hospitalizations. To date, 66 intensive care unit (ICU) admissions have been reported. A total of 42 ICU cases reported at least one underlying condition or comorbidity. The median age of patients admitted to the ICU was 68 years. Approximately 45 deaths have been reported this season, the majority in adults aged 65+. The median age of reported deaths was 85 years.

**Figure 8 - Cumulative numbers of adult hospitalizations ( $\geq 20$  years of age) with influenza by type and age-group reported by CIRN, Canada, 2016-17, Week 09**



**Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza by age-group ( $\geq 20$  years of age) reported by CIRN, Canada 2016-17, Week 09**



The number of hospitalizations reported through CIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

## Influenza Strain Characterizations

During the 2016-17 influenza season, the National Microbiology Laboratory (NML) has characterized 1,075 influenza viruses [969 A(H3N2), 25 A(H1N1), 81 influenza B]. All but one influenza A virus (n=993) and 32 influenza B viruses characterized were antigenically or genetically similar to the vaccine strains included in both the trivalent and quadrivalent vaccines. Forty-nine influenza B viruses were similar to the strain which is only included in the quadrivalent vaccine.

The World Health Organization (WHO) has released the recommended composition of the influenza vaccine for use in the 2017-2018 northern hemisphere influenza season. Trivalent vaccines are recommended to contain: 1) an A/Michigan/45/2015 (H1N1)pdm09-like virus; 2) an A/Hong Kong/4801/2014 (H3N2)-like virus; and 3) a B/Brisbane/60/2008-like virus (Victoria lineage). Quadrivalent vaccines are recommended to contain the above three viruses and a B/Phuket/3073/2013-like virus (Yamagata lineage).

**Table 3 – Influenza strain characterizations, Canada, 2016-17, Week 09**

Strain Characterization Results <sup>1</sup>	Count	Description
<b>Influenza A (H3N2)</b>		
Antigenically A/Hong Kong/4801/2014-like	285	Viruses antigenically similar to A/Hong Kong/4801/2014, the A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.
Genetically <sup>2</sup> A/Hong Kong/4801/2014-like	683	Viruses belonging to genetic group 3C.2a. A/Hong Kong/4801/2014-like virus belongs to genetic group 3C.2a and is the influenza A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.  Additionally, genetic characterization of the 285 influenza A (H3N2) viruses that underwent HI testing determined that 240 viruses belonged to genetic group 3C.2a and 38 viruses belonged to genetic group 3C.3a. Sequencing is pending for the remaining 7 isolates. The majority of viruses belonging to genetic group 3C.3a are inhibited by antisera raised against A/Hong Kong/4801/2014 <sup>3</sup> .
Antigenically A/Indiana/10/2011-like <sup>4</sup>	1	Viruses antigenically similar to A/Indiana/10/2011, a candidate H3N2v vaccine virus.
<b>Influenza A (H1N1)</b>		
A/California/7/2009-like	25	Viruses antigenically similar to A/California/7/2009, the A(H1N1) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
<b>Influenza B</b>		
B/Brisbane/60/2008-like (Victoria lineage)	32	Viruses antigenically similar to B/Brisbane/60/2008, the influenza B component of the 2016-17 Northern Hemisphere's <b>trivalent</b> and <b>quadrivalent</b> influenza vaccine.
B/Phuket/3073/2013-like (Yamagata lineage)	49	Viruses antigenically similar to B/Phuket/3073/2013, the additional influenza B component of the 2016-17 Northern Hemisphere <b>quadrivalent</b> influenza vaccine.

<sup>1</sup>The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#).

<sup>2</sup>Determined by sequence analysis

<sup>3</sup>[WHO](#) - Recommended composition of the influenza virus vaccines for use in the 2016-17 northern hemisphere influenza season.

<sup>4</sup>Detected in epidemiological week 50. For more details, see [Week 50 report](#)

## Antiviral Resistance

During the 2016-17 season, the National Microbiology Laboratory (NML) has tested 562 influenza viruses for resistance to oseltamivir and zanamivir and 158 influenza viruses for resistance to amantadine. All but one influenza A(H3N2) virus were sensitive to oseltamivir and all viruses were sensitive to zanamivir. All 158 influenza A viruses were resistant to amantadine (Table 4).

**Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2016-17, Week 09**

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	548	1 (0.2%)	548	0 (0%)	148	148 (100%)
A (H3N2v)	1	0 (0%)	1	0 (0%)	1	1 (100%)
A (H1N1)	22	0 (0%)	21	0 (0%)	20	20 (100%)
B	68	0 (0%)	69	0 (0%)	NA <sup>1</sup>	NA <sup>1</sup>
<b>TOTAL</b>	<b>639</b>	<b>1 (0.2%)</b>	<b>639</b>	<b>0 (0%)</b>	<b>169</b>	<b>169 (100%)</b>

<sup>1</sup>NA: Not Applicable

## Provincial and International Influenza Reports

- [World Health Organization influenza update](#)
- [World Health Organization FluNet](#)
- [WHO Influenza at the human-animal interface](#)
- [Centers for Disease Control and Prevention seasonal influenza report](#)
- [European Centre for Disease Prevention and Control - epidemiological data](#)
- [South Africa Influenza surveillance report](#)
- [New Zealand Public Health Surveillance](#)
- [Australia Influenza Report](#)
- [Pan-American Health Organization Influenza Situation Report](#)
- [Alberta Health – Influenza Surveillance Report](#)
- [BC - Centre for Disease Control \(BCCDC\) - Influenza Surveillance](#)
- [New Brunswick – Influenza Surveillance Reports](#)
- [Newfoundland and Labrador – Surveillance and Disease Reports](#)
- [Nova Scotia - Flu Information](#)
- [Public Health Ontario – Ontario Respiratory Pathogen Bulletin](#)
- [Manitoba – Epidemiology and Surveillance – Influenza Reports](#)
- [Saskatchewan – influenza Reports](#)
- [PEI – Influenza Summary](#)

## **FluWatch Definitions for the 2016-2017 Season**

**Abbreviations:** Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

**Influenza-like-illness (ILI):** Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

### **ILI/Influenza outbreaks**

**Schools:** Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

**Hospitals and residential institutions:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

**Workplace:** Greater than 10% absenteeism on any day which is most likely due to ILI.

**Other settings:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

*Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.*

### **Influenza/ILI Activity Levels**

**1 = No activity:** no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

**2 = Sporadic:** sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

**3 = Localized:** (1) evidence of increased ILI\* ;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

**4 = Widespread:** (1) evidence of increased ILI\*;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

*Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.*

*\* More than just sporadic as determined by the provincial/territorial epidemiologist.*

*† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.*

*We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.*

This [report](#) is available on the Government of Canada Influenza webpage. Ce rapport est disponible dans les deux langues officielles.