

March 12 to March 18, 2017 (Week 11)


Overall Summary

- Overall, the slow decline in influenza activity in Canada has continued in week 11. However, many parts of Canada, particularly the Eastern and Atlantic regions are still reporting elevated activity in week 11.
- In week 11, the number of laboratory detections, outbreaks and the number of geographic regions with influenza activity, decreased from the previous week.
- Although adult sentinel hospitalizations decreased from the previous week, the number of hospitalizations and deaths reported by participating provinces and territories increased.
- Influenza B detections and outbreaks in Canada are slowly increasing.
- Although declining for most indicators, influenza A(H3N2) continues to be the most common subtype of influenza affecting Canadians.
- The majority of laboratory detections, hospitalizations and deaths have been among adults aged 65+ years.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In week 11, one region in NL, reported no influenza or influenza-like illness activity. Sporadic influenza activity was reported in 29 regions across 11 provinces and territories. Localized activity was reported in 21 regions across nine provinces. No regions reported any widespread activity in week 11. For more details on a specific region, click on the map.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 11




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

Laboratory Confirmed Influenza Detections

In week 11, the number (1,197) and the percentage of tests positive for influenza (16%) decreased from the previous week. Peak influenza detections occurred in week 02 at 27%. Although declining, influenza A continues to account for the majority of detections; however, influenza B detections have been steadily increasing for the past few weeks. Influenza B activity is very low compared to the same time period in the previous two seasons. For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada (PHAC) website.


Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2016-17, Week 11



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

To date this season, 32,836 laboratory confirmed influenza detections have been reported, of which 96% have been influenza A. Influenza A(H3N2) is the most common subtype detected. For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2016-17, Week 11



To date, detailed information on age and type/subtype has been received for 23,013 laboratory-confirmed influenza cases (Table 1). Among cases with reported age and type/subtype information, adults aged 65+ accounted for almost half of the reported influenza cases. Among cases of influenza A(H3N2), adults aged 65+ represented 49% of cases, followed by adults aged 20-64 (34% of cases). In the previous influenza A(H3N2)-predominant season in 2014-15, adults aged 65+ represented 58% of cases and adults aged 20-64 represented 27% of cases.

Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting¹, Canada, 2016-17, Week 11

Age groups (years)	Week (March 12 to March 18, 2017)					Cumulative (August 28, 2016 to March 18, 2017)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) ³		A Total	A(H1) pdm09	A(H3)	A (UnS) ³		Total	#
0-4	>87	<5	10	77	7	2068	14	799	1255	102	2170	9%
5-19	>24	0	<5	24	21	2076	13	1038	1025	172	2248	10%
20-44	64	0	7	57	32	3233	27	1751	1455	177	3410	15%
45-64	85	0	17	68	19	3595	23	1871	1701	210	3805	17%
65+	>321	<5	64	257	43	11063	13	5204	5846	317	11380	49%
Total	586	<5	>98	483	122	22035	90	10663	11282	978	23013	100%
Percentage²	83%	x%	x%	82%	17%	96%	0%	48%	51%	4%		

¹Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. Cumulative data include updates to previous weeks.

²Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

³UnS: unsubtyped: The specimen was typed as influenza A, but no result for subtyping was available.

x: Suppressed to prevent residual disclosure

Specimens from NT, YT, and NU are sent to reference laboratories in the provinces


Syndromic/Influenza-like Illness Surveillance

Healthcare Professionals Sentinel Syndromic Surveillance

In week 11, 1.7% of visits to healthcare professionals were due to influenza-like illness, compared to 1.8% in the previous week.

Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2016-17

Number of Sentinels Reporting Week 11: 101



Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.


Are you a primary healthcare practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel? Please visit our [Influenza Sentinel page](#) for more details.

Influenza Outbreak Surveillance

In week 11, 42 laboratory confirmed influenza outbreaks were reported (three less than the previous week). Among the reported outbreaks: 21 were in long-term care (LTC) facilities, seven in hospitals and 12 in institutional or community (other) settings. Of the outbreaks with known strains or subtypes, 11 were due to influenza A(H3N2), eight were due to influenza A(UnS) and seven outbreaks were due to influenza B. An additional two outbreaks due to ILI were reported in a school.

To date this season, 991 outbreaks have been reported and the majority (66%) have occurred in LTC facilities. Compared to the same period in the most recent previous A(H3N2) predominant season (2014-15), 1,552 outbreaks were reported, of which 74% occurred in LTC facilities.

Figure 5 – Number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2016-17, Week 11



¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

Provincial/Territorial Influenza Hospitalizations and Deaths

In week 11, 261 influenza-associated hospitalizations were reported by participating provinces and territories, up from 254 reported in the previous week. Influenza A accounted for 88% of hospitalizations. The weekly percentage of influenza B associated hospitalizations has been steadily increasing for the past few weeks (from 1.2% in week 02 to 11.5% in week 11). The largest proportion of hospitalizations were among adults aged 65+ years (71%). A total of ten intensive care unit (ICU) admissions were reported in week 11. An increase in the number of deaths was reported in week 11, with 35 deaths reported. All deaths were reported in adults aged 65+ years.

To date this season, 5,139 hospitalizations have been reported, of which 97% were due to influenza A. Among cases for which the subtype of influenza A was reported, almost all (2770/2787) were influenza A(H3N2). Adults 65+ accounted for 69% of the hospitalizations. A total of 193 ICU admissions and 277 deaths have been reported. The majority of deaths was reported in adults aged 65+ years.

Table 2 – Cumulative number of hospitalizations, ICU admissions and deaths by age and influenza type reported by participating provinces and territories, Canada, 2016-17, Week 11

Age Groups (years)	Cumulative (August 28, 2016 to March 18, 2017)						
	Hospitalizations			ICU Admissions		Deaths	
	Influenza A Total	Influenza B Total	Total [# (%)]	Influenza A and B Total	%	Influenza A and B Total	%
0-4	395	17	412 (8%)	11	6%	<5	x%
5-19	217	21	238 (5%)	13	7%	<5	x%
20-44	263	8	271 (5%)	18	9%	<5	x%
45-64	666	30	696 (14%)	52	27%	32	12%
65+	3433	89	3522 (68%)	99	51%	238	86%
Total	4974	165	5139 (100%)	193	100%	277	100%

x: Suppressed to prevent residual disclosure

*Note: Influenza-associated hospitalizations are not reported to PHAC by BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

Sentinel Hospital Influenza Surveillance

Pediatric Influenza Hospitalizations and Deaths

In week 11, 17 laboratory-confirmed influenza-associated pediatric (≤ 16 years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. All but three cases were due to influenza A. The number of weekly hospitalizations reported since week 05 have been below the six year average for the same time period (Figure 7).

To date this season, 458 laboratory-confirmed influenza-associated pediatric hospitalizations were reported by the IMPACT network. Children aged 0-23 months accounted for approximately 39% of hospitalizations. Influenza A accounted for 91% (n=419) of the reported hospitalizations, of which 34% (n=141) were influenza A(H3N2) and the remainder were A(UnS). Additionally, 74 intensive care unit (ICU) admissions have been reported. Children aged 10-16 years accounted for 30% of ICU cases followed by children aged 0-23 months (27%). A total of 47 ICU cases reported at least one underlying condition or comorbidity. Less than five deaths have been reported this season.

Figure 6 – Cumulative numbers of pediatric hospitalizations (≤ 16 years of age) with influenza by age-group reported by the IMPACT network, Canada, 2016-17, Week 11





Figure 7 – Number of pediatric hospitalizations (≤ 16 years of age) with influenza reported by the IMPACT network, by week, Canada, 2016-17, Week 11



The shaded area represents the maximum and minimum number of cases reported by week from seasons 2010-11 to 2015-16

The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Adult Influenza Hospitalizations and Deaths

In week 11, 42 laboratory-confirmed influenza-associated adult (≥ 20 years of age) hospitalizations were reported by the Canadian Immunization Research Network (CIRN). All but nine cases were due to influenza A and the majority of cases (69%) occurred in adults aged 65+. The number of hospitalizations due to influenza B has been increasing since week 05.

To date this season, 1,222 laboratory-confirmed influenza-associated adult (≥ 20 years of age) hospitalizations have been reported by CIRN. Influenza A accounted for 98% of hospitalizations. Adults aged 65+ accounted for 78% of hospitalizations. To date, 81 intensive care unit (ICU) admissions have been reported. A total of 55 ICU cases reported at least one underlying condition or comorbidity. The median age of patients admitted to the ICU was 69 years. Approximately 51 deaths have been reported this season, the majority in adults aged 65+. The median age of reported deaths was 85 years.

Figure 8 - Cumulative numbers of adult hospitalizations (≥ 20 years of age) with influenza by type and age-group reported by CIRN, Canada, 2016-17, Week 11





Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza by age-group (≥ 20 years of age) reported by CIRN, Canada 2016-17, Week 11



The number of hospitalizations reported through CIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Influenza Strain Characterizations

During the 2016-17 influenza season, the National Microbiology Laboratory (NML) has characterized 1,420 influenza viruses [1272 A(H3N2), 28 A(H1N1), 120 influenza B]. All but one influenza A virus (n=1271) and 40 influenza B viruses characterized were antigenically or genetically similar to the vaccine strains included in both the trivalent and quadrivalent vaccines. Eighty influenza B viruses were similar to the strain which is only included in the quadrivalent vaccine.

The World Health Organization (WHO) has released the recommended composition of the influenza vaccine for use in the 2017-2018 northern hemisphere influenza season. Trivalent vaccines are recommended to contain: 1) an A/Michigan/45/2015 (H1N1)pdm09-like virus; 2) an A/Hong Kong/4801/2014 (H3N2)-like virus; and 3) a B/Brisbane/60/2008-like virus (Victoria lineage). Quadrivalent vaccines are recommended to contain the above three viruses and a B/Phuket/3073/2013-like virus (Yamagata lineage).

Table 3 – Influenza strain characterizations, Canada, 2016-17, Week 11

Strain Characterization Results ¹	Count	Description
Influenza A (H3N2)		
Antigenically A/Hong Kong/4801/2014-like	319	Viruses antigenically similar to A/Hong Kong/4801/2014, the A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.
Genetically ² A/Hong Kong/4801/2014-like	952	Viruses belonging to genetic group 3C.2a. A/Hong Kong/4801/2014-like virus belongs to genetic group 3C.2a and is the influenza A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine. Additionally, genetic characterization of the 319 influenza A (H3N2) viruses that underwent HI testing determined that 272 viruses belonged to genetic group 3C.2a and 47 viruses belonged to genetic group 3C.3a. Sequencing is pending for the remaining one isolate. The majority of viruses belonging to genetic group 3C.3a are inhibited by antisera raised against A/Hong Kong/4801/2014 ³ .
Antigenically A/Indiana/10/2011-like ⁴	1	Viruses antigenically similar to A/Indiana/10/2011, a candidate H3N2v vaccine virus.
Influenza A (H1N1)		
A/California/7/2009-like	28	Viruses antigenically similar to A/California/7/2009, the A(H1N1) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
Influenza B		
B/Brisbane/60/2008-like (Victoria lineage)	40	Viruses antigenically similar to B/Brisbane/60/2008, the influenza B component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
B/Phuket/3073/2013-like (Yamagata lineage)	80	Viruses antigenically similar to B/Phuket/3073/2013, the additional influenza B component of the 2016-17 Northern Hemisphere quadrivalent influenza vaccine.

¹The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#).

²Determined by sequence analysis

³[WHO](#) - Recommended composition of the influenza virus vaccines for use in the 2016-17 northern hemisphere influenza season.

⁴Detected in epidemiological week 50. For more details, see [Week 50 report](#)

Antiviral Resistance

During the 2016-17 season, the National Microbiology Laboratory (NML) has tested 760 influenza viruses for resistance to oseltamivir and zanamivir and 180 influenza viruses for resistance to amantadine. All but one influenza A(H3N2) virus were sensitive to oseltamivir and all viruses were sensitive to zanamivir. All 180 influenza A viruses were resistant to amantadine (Table 4).

Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2016-17, Week 11

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	635	1 (0.2%)	635	0 (0%)	155	155 (100%)
A (H3N2v)	1	0 (0%)	1	0 (0%)	1	1 (100%)
A (H1N1)	25	0 (0%)	24	0 (0%)	24	24 (100%)
B	99	0 (0%)	100	0 (0%)	NA ¹	NA ¹
TOTAL	760	1 (0.1%)	760	0 (0%)	180	180 (100%)

¹NA: Not Applicable

Provincial and International Influenza Reports

- [World Health Organization influenza update](#)
- [World Health Organization FluNet](#)
- [WHO Influenza at the human-animal interface](#)
- [Centers for Disease Control and Prevention seasonal influenza report](#)
- [European Centre for Disease Prevention and Control - epidemiological data](#)
- [South Africa Influenza surveillance report](#)
- [New Zealand Public Health Surveillance](#)
- [Australia Influenza Report](#)
- [Pan-American Health Organization Influenza Situation Report](#)
- [Alberta Health – Influenza Surveillance Report](#)
- [BC - Centre for Disease Control \(BCCDC\) - Influenza Surveillance](#)
- [New Brunswick – Influenza Surveillance Reports](#)
- [Newfoundland and Labrador – Surveillance and Disease Reports](#)
- [Nova Scotia - Flu Information](#)
- [Public Health Ontario – Ontario Respiratory Pathogen Bulletin](#)
- [Manitoba – Epidemiology and Surveillance – Influenza Reports](#)
- [Saskatchewan – influenza Reports](#)
- [PEI – Influenza Summary](#)

FluWatch Definitions for the 2016-2017 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This [report](#) is available on the Government of Canada Influenza webpage. Ce rapport est disponible dans les deux langues officielles.