

February 22 to February 28, 2015 (week 08)

## Overall Summary


- In week 08, all influenza indicators except the influenza-like illness consultation rate remained similar to, or declined from the previous week.
- Elevated influenza activity was mostly reported in the Central and Atlantic provinces and in a few regions in the Western provinces.
- Influenza B detections continues to increase steadily, particularly in the Prairies and in Quebec. This increase in influenza B is expected as influenza B often shows up later in the flu season.
- A(H3N2) continues to be the most common influenza virus this season and seniors continue to have the highest number of positive laboratory detections, hospitalizations and deaths.
- Evidence from the National Microbiology Laboratory (NML) indicates that this year's vaccine will continue to provide protection against the circulating A(H1N1) and B strains.

Are you a primary health care practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel for the 2014-15 influenza season? Contact us at [FluWatch@phac-aspc.gc.ca](mailto:FluWatch@phac-aspc.gc.ca)

## Influenza/ILI Activity (geographic spread)

In week 08, two regions in Quebec reported widespread activity. Twenty-three regions reported localized activity: BC(2), AB(2), MB, ON(7), QC(2), NB(4), NS(3), PE and NL. Twenty-five regions reported sporadic activity: in YK, NU, NT(2), BC(3), AB(3), SK(3), MB(3), QC(2), NB(3), NS(3), and NL. No activity was reported in seven regions: NU, MB, NS(3) and NL(2). Overall when compared to the previous week, there was a decrease in widespread activity. The majority of widespread and localized activity was reported in the Central and Atlantic provinces; however, more regions reported localised activity compared to the previous week.

Figure 1. Map of overall influenza/ILI activity level by province and territory, Canada, Week 08




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available on the [FluWatch website](http://FluWatch website).

## Influenza and Other Respiratory Virus Detections

In week 08, the number of positive influenza tests (1,242) and the percentage positive for influenza A (11.1%) continued to decline from the previous week (figure 2). The percentage of positive influenza B tests continued to increase and was 5.9% in week 08 (Figure 2). Influenza B detections were greater than influenza A detections in QC and AB; accounting for 57% and 63% of influenza detections respectively. To date, 93% of influenza detections have been influenza A, and 99.5% of those subtyped have been A(H3N2) (Table 1). To date this season, detailed information on age and type/subtype has been received for 30,350 cases. A significantly greater proportion of laboratory detections of influenza have been reported in adults  $\geq 65$  years of age (61%) this season (Table 2) compared to the 2013-14 season when only 16.3% of cases were in adults  $\geq 65$  years of age.


**Figure 2. Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2014-15**



In week 08, detections of all respiratory viruses except parainfluenza and human metapneumovirus (hMPV) decreased from the previous week (Figure 3). Detections of respiratory syncytial virus (RSV) in week 08 were greater than the detections of influenza A with 867 detections (vs 818 detections of influenza A). In recent weeks, weekly detections of adenovirus, coronavirus, rhinovirus and parainfluenza have been greater than those reported in each of the past three seasons. Weekly detections of hMPV this season have been lower compared to the previous three seasons.

For more details, see the weekly [Respiratory Virus Detections in Canada Report](#).

**Figure 3. Number of positive laboratory tests for other respiratory viruses by report week, Canada, 2014-15**



RSV: Respiratory syncytial virus; hMPV: Human metapneumovirus

**Table 1. Weekly and cumulative numbers of positive influenza specimens by type, subtype and province, Canada, 2014-15**

Reporting provinces <sup>1</sup>	Weekly (February 22 to February 28, 2015)					Cumulative (August 24, 2014 to February 28, 2015)				
	Influenza A				B	Influenza A				B
	A Total	A(H1)pdm09	A(H3)	A(UnS)	B Total	A Total	A(H1)pdm09	A(H3)	A(UnS)	B Total
BC	36	1	28	7	22	3408	24	2550	834	176
AB	23	0	15	8	39	3621	10	3455	156	446
SK	17	0	0	11	6	1293	0	0	1293	51
MB	12	0	3	9	5	1108	0	382	726	43
ON	341	7	113	221	48	10245	25	4357	5863	236
QC	200	0	0	200	260	10929	4	422	10503	1232
NB	145	0	26	119	25	864	0	128	736	75
NS	23	0	0	23	15	401	0	123	278	60
PE	6	0	6	0	4	111	1	108	2	6
NL	15	0	0	15	0	579	0	53	526	4
<b>Canada</b>	<b>818</b>	<b>8</b>	<b>191</b>	<b>613</b>	<b>424</b>	<b>32559</b>	<b>64</b>	<b>11578</b>	<b>20917</b>	<b>2329</b>
<b>Percentage<sup>2</sup></b>	65.9%	1.0%	23.3%	74.9%	34.1%	93.3%	0.2%	35.6%	64.2%	6.7%

**Table 2. Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting<sup>3</sup>, Canada, 2014-15**

Age groups (years)	Weekly (February 22 to February 28, 2015)					Cumulative (August 24, 2014 to February 28, 2015)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1)pdm09	A(H3)	A (UnS)	Total	A Total	A(H1)pdm09	A(H3)	A (UnS)	Total	#	%
<5	28	2	3	23	21	1936	12	723	1201	162	2098	6.9%
5-19	13	4	4	5	35	1682	5	895	782	269	1951	6.4%
20-44	26	0	5	21	60	3251	13	1460	1778	369	3620	11.9%
45-64	53	0	9	44	85	3609	13	1428	2168	512	4121	13.6%
65+	245	0	43	202	136	17677	10	6543	11124	766	18443	60.8%
Unknown	2	0	2	0	0	115	0	97	18	2	117	0.4%
<b>Total</b>	<b>367</b>	<b>6</b>	<b>66</b>	<b>295</b>	<b>337</b>	<b>28270</b>	<b>53</b>	<b>11146</b>	<b>17071</b>	<b>2080</b>	<b>30350</b>	<b>100.0%</b>
<b>Percentage<sup>2</sup></b>	52.1%	1.6%	18.0%	80.4%	47.9%	93.1%	0.2%	39.4%	60.4%	6.9%		

<sup>1</sup> Specimens from NT, YT, and NU are sent to reference laboratories in other provinces. Cumulative data includes updates to previous weeks.

<sup>2</sup> Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

<sup>3</sup> Table 2 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. UnS: unsubtype: The specimen was typed as influenza A, but no result for subtyping was available.

## Antiviral Resistance

During the 2014-2015 influenza season, the NML has tested 764 influenza viruses for resistance to oseltamivir and 759 influenza viruses for resistance to zanamivir. All viruses were sensitive to zanamivir and one influenza A(H3N2) virus was resistant to oseltamivir. A total of 1042 influenza A viruses (99.9%) were resistant to amantadine (Table 3).

**Table 3. Antiviral resistance by influenza virus type and subtype, Canada, 2014-15**


Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	651	1	646	0	1038	1037 (99.9%)
A (H1N1)	3	0	3	0	4	4 (100%)
B	110	0	110	0	NA <sup>1</sup>	NA <sup>1</sup>
<b>TOTAL</b>	<b>764</b>	<b>1</b>	<b>759</b>	<b>0</b>	<b>1042</b>	<b>1041</b>

<sup>1</sup>NA: Not Applicable

## Influenza Strain Characterizations

During the 2014-2015 influenza season, the National Microbiology Laboratory (NML) has characterized 277 influenza viruses [125 A(H3N2), 3 A(H1N1) and 149 influenza B].

**Influenza A (H3N2):** When tested by hemagglutination inhibition (HI) assay (n=125), one virus was antigenically similar to A/Texas/50/2012, five showed reduced titers to A/Texas/50/2012 and 119 were antigenically similar to A/Switzerland/9715293/2013, which is the influenza A(H3N2) component recommended for the 2015 Southern Hemisphere influenza vaccine. Additionally, 827 A(H3N2) viruses were unable to be tested by HI assay; however, sequence analysis showed that 825 belonged to a genetic group that typically shows reduced titers to A/Texas/50/2012. **Influenza A(H1N1):** Three A(H1N1) viruses characterized were antigenically similar to A/California/7/2009. **Influenza B:** Of the 149 influenza B viruses characterized, 142 viruses were antigenically similar to B/Massachusetts/2/2012, three viruses showed reduced titers against B/Massachusetts/2/2012, and four were B/Brisbane/60/2008-like (Figure 4).



**Figure 4. Influenza strain characterizations, Canada, 2014-15, N = 277**

The NML receives a proportion of the number of influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Characterization data reflect the results of haemagglutination inhibition (HAI) testing compared to the reference influenza strains recommended by [WHO](#).


The recommended components for the 2014-2015 northern hemisphere trivalent influenza vaccine include: an A/California/7/2009(H1N1)pdm09-like virus, an A/Texas/50/2012 (H3N2)-like virus, and a B/Massachusetts/2/2012-like virus (Yamagata lineage). For quadrivalent vaccines, the addition of a B/Brisbane/60/2008-like virus is recommended.

The WHO has released the recommended composition of the influenza vaccine for the northern hemisphere for the 2015-2016 season. Trivalent vaccines are recommended to contain 1) an A/California/7/2009 (H1N1)pdm09-like virus 2) an A/Switzerland/9715293/2013 (H3N2)-like virus, and 3) an B/Phuket/3073/2013-like virus (Yamagata lineage). Quadrivalent vaccines are recommended to additionally contain a B/Brisbane/60/2008-like virus (Victoria lineage).

## Influenza-like Illness Consultation Rate

The national influenza-like-illness (ILI) consultation rate increased to 52.0 consultations per 1,000, which is above expected levels for week 08 (Figure 5). The rate was highest among the 5 to 19 years of age group (67.7 consultations per 1,000) and lowest among the adults  $\geq 65$  years of age (34.6 consultations per 1,000)

**Figure 5. Influenza-like-illness (ILI) consultation rates by report week, compared to the 1996-97 through to 2012-13 seasons (with pandemic data suppressed), Canada, 2014-15**




No data available for mean rate for weeks 19 to 39 for the 1996-1997 through 2002-2003 seasons. Delays in the reporting of data may cause data to change retrospectively. The calculation of the average ILI consultation rate over 17 seasons was aligned with influenza activity in each season. In BC, AB, and SK, data is compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.

## Influenza Outbreak Surveillance

In week 08, 31 new outbreaks of influenza were reported, a decrease from the previous week. The majority of the outbreaks occurred in the Central and Atlantic provinces. Twenty-four outbreaks were reported in long-term care facilities (LTCF) and seven in institutional or community settings (Figure 6). Among the outbreaks in which the influenza subtype was known, two LTCF outbreaks were associated with A(H3N2). To date this season, 1,039 outbreaks in LTCFs have been reported and the majority of those with known subtypes were attributable to A(H3N2). There have been a higher number of reported influenza outbreaks to date this season compared to the same period in previous seasons.

**Figure 6. Overall number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2014-2015**




<sup>1</sup>All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of the report.

## Pharmacy Surveillance

During week 08, the proportion of prescriptions for antivirals decreased to 157.8 antiviral prescriptions per 100,000 total prescriptions (from 221.8 per 100,000). The rate for antivirals since week 48 has been higher than the previous three seasons (Figure 7). The rate in all age groups decreased in week 08. The rate was highest among seniors at 260.3 per 100,000 total prescriptions and lowest among infants at 45.2 per 100,000 total prescriptions.

**Figure 7 – Proportion of prescription sales for influenza antivirals by age-group and week, Canada, 2014-15**



Note: Pharmacy sales data are provided to the Public Health Agency of Canada by Rx Canada Inc. and sourced from major retail drug chains representing over 2,500 stores nationwide (excluding Nunavut) in 85% of Health Regions. Data provided include the number of new antiviral prescriptions (for Tamiflu and Relenza) and the total number of new prescriptions dispensed by Province/Territory and age group. Age-groups: Infant: 0-2y, Child: 2-18y; Adult: 19-64y, Senior: ≥65y

## Sentinel Hospital Influenza Surveillance

### Paediatric Influenza Hospitalizations and Deaths (IMPACT)

In week 08, nine laboratory-confirmed influenza-associated paediatric ( $\leq 16$  years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network: four cases of influenza A and five cases of influenza B (Figure 8a). A greater proportion of cases have been reported with influenza B in recent weeks, following the trend in laboratory detections. Among the reported cases, four (44%) were  $< 2$  years of age, three (44%) were 2 to 9 years of age and one (11%) was 10-16 years of age. One case was admitted to the ICU.

To date this season, 541 hospitalizations have been reported by the IMPACT network, 486 (90%) of which were cases of influenza A. Among cases for which the influenza A subtype was reported, 99% (157/159) were A(H3N2) (Table 4). To date, 66 cases were admitted to the ICU, of which 36 (55%) were 2 to 9 years of age (Figure 9a). A total of 38 ICU cases reported to have at least one underlying condition or comorbidity. Three deaths have been reported.

Note: The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated paediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

### Adult Influenza Hospitalizations and Deaths (PCIRN)

In week 08, 67 laboratory-confirmed influenza-associated adult ( $\geq 16$  years of age) hospitalizations were reported by the PHAC/CIHR Influenza Research Network (PCIRN) Serious Outcomes Surveillance (SOS) network. Among the cases in week 08, 50 cases (75%) were in adults over the age of 65 and 53 cases (79%) had influenza A (Figure 8b).

To date this season, 1,810 cases have been reported; 1,741 (96%) with influenza A. The majority of cases (82%) were among adults  $\geq 65$  years of age (Table 5). One hundred and thirty-six ICU admissions have been reported and 103 cases were adults  $\geq 65$  years of age. A total of 97 ICU cases (71%) reported to have at least one underlying condition or comorbidity. Of the 94 ICU cases with known immunization status, 33 (35%) reported not having been vaccinated this season. Ninety-three deaths have been reported, 85 (91%) of the deaths were adults  $> 65$  years of age (Figure 9B).

Note: The number of hospitalizations reported through PCIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

**Table 4 – Cumulative numbers of paediatric hospitalizations with influenza reported by the IMPACT network, Canada, 2014-15**

Age groups	Cumulative (24 Aug. 2014 to 28 Feb. 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>2</sup>	Total	# (%)
0-5m	76	0	16	60	3	79 (14.6%)
6-23m	107	1	35	71	20	127 (23.5%)
2-4y	120	1	39	80	12	132 (24.4%)
5-9y	125	0	44	81	12	137 (25.3%)
10-16y	58	0	23	35	8	66 (12.2%)
<b>Total</b>	486	2	157	327	55	541
<b>%<sup>1</sup></b>	89.8%	0.4%	32.3%	67.3%	10.2%	100.0%


**Table 5 – Cumulative numbers of adult hospitalizations with influenza reported by the PCIRN-SOS network, Canada, 2014-15**

Age groups (years)	Cumulative (15 Nov. 2014 to 28 Feb. 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A (UnS)	Total	# (%)
16-20	5	0	1	4	0	5 (%)
20-44	97	1	40	56	4	101 (6%)
45-64	195	0	79	116	19	214 (12%)
65+	1444	3	576	865	46	1490 (82%)
<b>Total</b>	1741	4	696	1041	69	1810
<b>%</b>	96%	0%	40%	60%	4%	100%


<sup>1</sup> Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections. UnS: unsubtype: The specimen was typed as influenza A, but no result for subtyping was available.

**Figure 8 – Number of cases of influenza reported by sentinel hospital networks, by week, Canada, 2014-15**

**A) Paediatric hospitalizations (≤16 years of age, IMPACT)**



**B) Adult hospitalizations (≥16 year of age, PCIRN-SOS)**




Note: Data for week 46 is based on data collected for 1 day only and do not represent the number of hospitalizations for the entire week.

**Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age-group, Canada, 2014-15**

**A) Paediatric hospitalizations (≤16 years of age, IMPACT)**



## B) Adult hospitalizations (≥16 year of age, PCIRN-SOS)



## Provincial/Territorial Influenza Hospitalizations and Deaths

In week 08, 213 laboratory-confirmed influenza-associated hospitalizations were reported from participating provinces and territories\* which is less than the number reported in week 07 (n=276). Of the 213 hospitalizations, all but 16 were due to influenza A, and 73% were in patients ≥65 years of age. Since the start of the 2014-15 season, 5,493 hospitalizations have been reported; 5,312 (97%) with influenza A. Among cases for which the subtype of influenza A was reported, 99.5% were A(H3N2). The majority of cases (71%) were ≥65 years of age (Table 6). A total of 274 ICU admissions have been reported to date: 54% (n=149) were in adults ≥65 years of age and 30% (n=83) were in adults 20-64 years. A total of 389 deaths have been reported since the start of the season: three children <5 years of age, one child 5-19 years, 32 adults 20-64 years, and 353 adults ≥65 years of age. Adults 65 years of age or older represent 91% of all deaths reported this season. Detailed clinical information (e.g. underlying medical conditions) is not known for these cases.

\* Note: Influenza-associated hospitalizations are not reported to PHAC by the following Provinces and Territory: BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by Saskatchewan. ICU admissions are not distinguished among hospital admissions reported from Ontario. Data may also include cases reported by the IMPACT and PCIRN networks. The number of new influenza-associated hospitalizations and deaths reported for the current week may include cases from Ontario that occurred in previous weeks, as a result of retrospective updates to the cumulative total. It is important to note that the hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

**Table 6 – Cumulative number of hospitalizations with influenza reported by the participating provinces and territories, Canada, 2014-15**

Age groups (years)	Cumulative (24 Aug. 2014 to 28 Feb. 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A (UnS)	Total	# (%)
0-4	363	2	132	229	8	371 (7%)
5-19	218	0	107	111	16	234 (4%)
20-44	263	3	133	127	18	281 (5%)
45-64	581	5	279	297	35	616 (11%)
65+	3832	1	1765	2066	94	3926 (71%)
Unknown	55	1	51	3	10	65 (1%)
<b>Total</b>	<b>5312</b>	<b>12</b>	<b>2467</b>	<b>2833</b>	<b>181</b>	<b>5493</b>
<b>Percentage<sup>1</sup></b>	<b>96.7%</b>	<b>0.2%</b>	<b>46.4%</b>	<b>53.3%</b>	<b>3.3%</b>	<b>100.0%</b>

<sup>1</sup> Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.  
UnS: untyped: The specimen was typed as influenza A, but no result for subtyping was available.

See additional data on [Reported Influenza Hospitalizations and Deaths in Canada: 2009-10 to 2014-15](#) on the Public Health Agency of Canada website.

## Emerging Respiratory Pathogens

### Human Avian Influenza

Influenza A(H7N9): Since the last FluWatch report, no new laboratory-confirmed cases of human infection with avian influenza A(H7N9) virus were reported by the World Health Organization. Globally to March 6, 2015, the WHO reported a total of 572 laboratory-confirmed human cases with avian influenza A(H7N9) virus, including 204 deaths. Documents related to the public health risk of influenza A(H7N9), as well as guidance for health professionals and advice for the public is updated regularly on the following websites:

[PHAC – Avian influenza A\(H7N9\)](#)

[WHO – Avian Influenza A\(H7N9\)](#)

Influenza A(H5N6): Since the last FluWatch report, no new cases of human infection with avian influenza A (H5N6) virus from China has been reported by the World Health Organization. Globally to March 5, 2015, the WHO has been informed of a total of three cases of avian influenza A (H5N6) virus, including two deaths.

### Middle East Respiratory Syndrome Coronavirus (MERS-CoV)

Since the last FluWatch report, 10 new laboratory-confirmed cases of MERS-CoV have been reported by the World Health Organization. Globally, from September 2012 to March 6, 2015, the WHO has been informed of a total of 1,040 laboratory-confirmed cases of infection with MERS-CoV, including 383 deaths. All cases have either occurred in the Middle East or have had direct links to a primary case infected in the Middle East. The public health risk posed by MERS-CoV in Canada remains low (see the [PHAC Assessment of Public Health Risk](#)) and for the latest global risk assessment posted by the WHO on February 5, 2015: [WHO MERS-CoV](#)

Documents related to the public health risk of MERS-CoV, as well as guidance for health professionals and advice for the public is updated regularly on the following websites:

[PHAC – Middle East respiratory syndrome coronavirus \(MERS-CoV\)](#)

[WHO – Coronavirus infections](#)

### Avian Influenza A(H5)

Since the last FluWatch report, no new information has been provided by the CFIA. Further information on the outbreak is provided on the following CFIA website.

[CFIA - Notifiable Avian Influenza](#)

For the latest Travel Health Notice on Avian Influenza (H5N1) visit the following webpage: [PHAC – Travel Health Notice](#)

## International Influenza Reports

[World Health Organization influenza update](#)

[World Health Organization FluNet](#)

[WHO Influenza at the human-animal interface](#)

[Centers for Disease Control and Prevention seasonal influenza report](#)

[European Centre for Disease Prevention and Control - epidemiological data](#)

[South Africa Influenza surveillance report](#)

[New Zealand Public Health Surveillance](#)

[Australia Influenza Report](#)

[Pan-American Health Organization Influenza Situation Report](#)

## **FluWatch Definitions for the 2014-2015 Season**

**Abbreviations:** Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

**Influenza-like-illness (ILI):** Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

### **ILI/Influenza outbreaks**

**Schools:** Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

**Hospitals and residential institutions:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

**Workplace:** Greater than 10% absenteeism on any day which is most likely due to ILI.

**Other settings:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

*Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.*

### **Influenza/ILI Activity Levels**

**1 = No activity:** no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

**2 = Sporadic:** sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

**3 = Localized:** (1) evidence of increased ILI\* ;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

**4 = Widespread:** (1) evidence of increased ILI\*;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

*Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.*

*\* More than just sporadic as determined by the provincial/territorial epidemiologist.*

*† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.*

*We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.*

This report is available on the Public Health Agency website at the following address: <http://www.phac-aspc.gc.ca/fluwatch/index.html>.

Ce rapport est disponible dans les deux langues officielles.