

August 2 to August 15, 2015 (weeks 31 and 32)

Overall Summary

- Overall, there is little to no influenza activity in Canada; however, in week 32, there were low levels of influenza activity reported in regions of British Columbia and Ontario.
- There was one influenza outbreak in a long term care facility reported in week 32.
- Rhinovirus was the most commonly detected respiratory virus in weeks 31 and 32
- As of week 32, 8,008 hospitalizations and 604 deaths have been reported from participating regions, which is more than were reported last year at this time (5,442 hospitalizations and 342 deaths).

Are you a primary health care practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel for the 2015-16 influenza season? Contact us at FluWatch@phac-aspc.gc.ca

Influenza/ILI Activity (geographic spread)

In week 32, sporadic and/or localized activity was reported BC and ON. Overall, there is low influenza/ILI activity in Canada.

Figure 1. Map of overall influenza/ILI activity level by province and territory, Canada, Week 32




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available on the FluWatch website.

Influenza and Other Respiratory Virus Detections

Influenza detections are at inter-seasonal levels. For the past six weeks, less than 1% of tests were positive for influenza (Figure 2). Most jurisdictions (with the exception of BC, AB and ON) reported no influenza detections in week 32 (Table 1). To date this season, detailed information on age and type/subtype has been received for 37,468 cases (Table 2). Adults ≥ 65 years of age have predominantly been affected by influenza A, accounting for 62% of influenza A detections.


Figure 2. Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2014-15



Detections for all other respiratory viruses have continued to decline and have been at inter-seasonal levels for the past weeks (figure 3). In weeks 31 and 32, rhinovirus was the predominant virus among other respiratory viruses.

For more details, see the weekly [Respiratory Virus Detections in Canada Report](#).

Figure 3. Number of positive laboratory tests for other respiratory viruses by report week, Canada, 2014-15



RSV: Respiratory syncytial virus; hMPV: Human metapneumovirus

Table 1. Weekly and cumulative numbers of positive influenza specimens by type, subtype and province, Canada, 2014-15

Reporting provinces ¹	Weekly (August 2 to August 15, 2015)					Cumulative (August 24, 2014 to August 15, 2015)				
	Influenza A				B	Influenza A				B
	A Total	A(H1)pdm09	A(H3)	A(UnS)	B Total	A Total	A(H1)pdm09	A(H3)	A(UnS)	B Total
BC	2	0	2	0	0	3564	28	2661	875	507
AB	1	0	1	0	0	3716	14	3547	155	1007
SK	0	0	0	0	0	1320	0	841	479	412
MB	0	0	0	0	0	1124	1	390	733	228
ON	1	0	1	0	0	11219	53	4754	6412	1555
QC	0	0	0	0	0	11464	4	422	11038	3914
NB	0	0	0	0	0	1196	0	193	1003	536
NS	0	0	0	0	0	511	1	123	387	263
PE	0	0	0	0	0	131	1	128	2	109
NL	0	0	0	0	0	629	0	123	506	81
Canada	4	0	4	0	0	34874	102	13182	21590	8612
Percentage²	100.0%	0.0%	100.0%	0.0%	0.0%	80.2%	0.3%	37.8%	61.9%	19.8%

Table 2. Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting³, Canada, 2014-15

Age groups (years)	Weekly (August 2 to August 15, 2015)					Cumulative (August 24, 2014 to August 15, 2015)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1)pdm09	A(H3)	A (UnS)	Total	A Total	A(H1)pdm09	A(H3)	A (UnS)	Total	#	%
<5	0	0	0	0	0	2095	23	811	1261	570	2665	7.1%
5-19	0	0	0	0	0	1786	6	959	821	809	2595	6.9%
20-44	1	0	1	0	0	3466	18	1685	1763	1154	4620	12.3%
45-64	3	0	2	1	0	3897	23	1673	2201	1845	5742	15.3%
65+	0	0	0	0	0	18787	13	7318	11456	2932	21719	58.0%
Unknown	0	0	0	0	0	120	0	101	19	7	127	0.3%
Total	4	0	3	1	0	30151	83	12547	17521	7317	37468	100.0%
Percentage²	100.0%	0.0%	75.0%	25.0%	0.0%	80.5%	0.3%	41.6%	58.1%	19.5%		

¹ Specimens from NT, YT, and NU are sent to reference laboratories in other provinces. Cumulative data includes updates to previous weeks.

² Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

³ Table 2 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. UnS: unsubtype: The specimen was typed as influenza A, but no result for subtyping was available.

Antiviral Resistance

During the 2014-2015 influenza season, the NML has tested 1,925 influenza viruses for resistance to oseltamivir and 1,923 influenza viruses for resistance to zanamivir. All viruses were sensitive to zanamivir and one influenza A(H3N2) virus was resistant to oseltamivir. A total of 1,492 influenza A viruses (99.9%) were resistant to amantadine (Table 3).

Table 3. Antiviral resistance by influenza virus type and subtype, Canada, 2014-15

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	978	1	976	0	1468	1467 (99.9%)
A (H1N1)	25	0	25	0	25	25 (100%)
B	922	0	922	0	NA ¹	NA ¹
TOTAL	1925	1	1923	0	1493	1492

¹NA: Not Applicable

Influenza Strain Characterizations

During the 2014-2015 influenza season, the National Microbiology Laboratory (NML) has characterized 1,162 influenza viruses [216 A(H3N2), 24 A(H1N1) and 922 influenza B].

Influenza A (H3N2): When tested by hemagglutination inhibition (HI) assay (n=216), one virus was antigenically similar to A/Texas/50/2012, five showed reduced titers to A/Texas/50/2012 and 210 were antigenically similar to A/Switzerland/9715293/2013, which is the influenza A(H3N2) component recommended for the 2015 Southern Hemisphere influenza vaccine. Additionally, 1,224 A(H3N2) viruses were unable to be tested by HI assay; however, sequence analysis showed that 1,222 belonged to a genetic group that typically shows reduced titers to A/Texas/50/2012. **Influenza A(H1N1):** 24 A(H1N1) viruses characterized were antigenically similar to A/California/7/2009. **Influenza B:** Of the 922 influenza B viruses characterized, 811 viruses were antigenically similar to B/Massachusetts/2/2012, three viruses showed reduced titers against B/Massachusetts/2/2012, and 108 were B/Brisbane/60/2008-like (Figure 4).





Figure 4. Influenza strain characterizations, Canada, 2014-15, N = 1,162

The NML receives a proportion of the number of influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Characterization data reflect the results of haemagglutination inhibition (HAI) testing compared to the reference influenza strains recommended by [WHO](http://www.who.int).

Influenza-like Illness Consultation Rate

The national influenza-like-illness (ILI) consultation rate decreased from 10.4 consultations per 1,000 in week 30 to 7.5 per 1,000 in week 31 and 5.1 per 1,000 in week 32 (Figure 5). The rates for week 31 to 32 have been slightly above the expected range for this time of year.

Figure 5. Influenza-like-illness (ILI) consultation rates by report week, compared to the 1996-97 through to 2012-13 seasons (with pandemic data suppressed), Canada, 2014-15




No data available for mean rate for weeks 19 to 39 for the 1996-1997 through 2002-2003 seasons. Delays in the reporting of data may cause data to change retrospectively. The calculation of the average ILI consultation rate over 17 seasons was aligned with influenza activity in each season. In BC, AB, and SK, data is compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.

Influenza Outbreak Surveillance

In week 32, one new outbreak of influenza in a long term care facility (LTCF) was reported (Figure 6). To date this season, 1,281 outbreaks in LTCFs have been reported and the majority of those with known subtypes were attributable to A(H3N2). There have been a higher number of reported influenza outbreaks to date this season compared to the same period in previous seasons.

Figure 6. Overall number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2014-2015




¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of the report.

Pharmacy Surveillance

Pharmacy surveillance for sales of influenza antivirals has ended for the 2014-2015 influenza season (Figure 7).

Figure 7 – Proportion of prescription sales for influenza antivirals by age-group and week, Canada, 2014-15



Note: Pharmacy sales data are provided to the Public Health Agency of Canada by Rx Canada Inc. and sourced from major retail drug chains representing over 2,500 stores nationwide (excluding Nunavut) in 85% of Health Regions. Data provided include the number of new antiviral prescriptions (for Tamiflu and Relenza) and the total number of new prescriptions dispensed by Province/Territory and age group. Age-groups: Infant: 0-2y, Child: 2-18y; Adult: 19-64y, Senior: ≥65y

Sentinel Hospital Influenza Surveillance

Paediatric Influenza Hospitalizations and Deaths (IMPACT)

In weeks 31 and 32, no laboratory-confirmed influenza-associated paediatric (≤ 16 years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. No ICU admissions were reported.

To date this season, 714 hospitalizations have been reported by the IMPACT network, 511 (72%) of which were cases of influenza A. Among cases for which the influenza A subtype was reported, 98% (164/167) were A(H3N2) (Table 4). To date, 104 cases were admitted to the ICU, of which 58 (56%) were 2 to 9 years of age (Figure 9a). A total of 68 ICU cases reported to have at least one underlying condition or comorbidity. Five deaths have been reported.

Note: The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated paediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Adult Influenza Hospitalizations and Deaths (CIRN)

Surveillance has ended for the 2014-2015 influenza season.

This season, 2,228 cases have been reported; 1,912 (86%) with influenza A. The majority of cases (81%) were among adults ≥ 65 years of age (Table 5). One hundred and seventy two ICU admissions have been reported and 128 cases were adults ≥ 65 years of age. Among the 172 ICU admissions, 27 were due to influenza B (12 in adults 45 to 64 years of age and 15 in adults over the age of 65). A total of 123 ICU cases (72%) reported to have at least one underlying condition or comorbidity. Of the 123 ICU cases with known immunization status, 40 (33%) reported not having been vaccinated this season. One hundred and thirty-five deaths have been reported, 124 (92%) of the deaths were adults > 65 years of age (Figure 9B).

Note: The number of hospitalizations reported through PCIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Table 4 – Cumulative numbers of paediatric hospitalizations with influenza reported by the IMPACT network, Canada, 2014-15

Age groups	Cumulative (24 Aug. 2014 to 15 Aug. 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A (UnS)	Total	# (%)
0-5m	84	0	19	65	16	100 (14.0%)
6-23m	115	2	37	76	44	159 (22.3%)
2-4y	122	1	39	82	52	174 (24.4%)
5-9y	129	0	44	85	55	184 (25.8%)
10-16y	61	0	25	36	36	97 (13.6%)
Total	511	3	164	344	203	714
%¹	71.6%	0.6%	32.1%	67.3%	28.4%	100.0%


Table 5 – Cumulative numbers of adult hospitalizations with influenza reported by the PCIRN-SOS network, Canada, 2014-15

Age groups (years)	Cumulative (15 Nov. 2014 to 2 May 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A(UnS)	Total	# (%)
16-20	3	0	1	2	1	4 (0.2%)
20-44	106	1	56	49	16	122 (5%)
45-64	217	3	99	115	76	293 (13%)
65+	1586	4	760	822	223	1809 (81%)
Total	1912	8	916	988	316	2228
%	86%	0.4%	48%	52%	14%	100%


¹ Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections. UnS: untyped: The specimen was typed as influenza A, but no result for subtyping was available.

Figure 8 – Number of cases of influenza reported by sentinel hospital networks, by week, Canada, 2014-15

A) Paediatric hospitalizations (≤ 16 years of age, IMPACT)




B) Adult hospitalizations (≥ 16 year of age, PCIRN-SOS)




Note: Data for week 46 is based on data collected for 1 day only and do not represent the number of hospitalizations for the entire week.

Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age-group, Canada, 2014-15

A) Paediatric hospitalizations (≤ 16 years of age, IMPACT)



B) Adult hospitalizations (≥16 year of age, PCIRN-SOS)



Provincial/Territorial Influenza Hospitalizations and Deaths

In week 32, 34 laboratory-confirmed influenza-associated hospitalizations were reported from participating provinces and territories*. These hospitalizations likely represent case updates due to delayed reporting. Of the 34 hospitalizations, 10 (29%) were due to influenza A and 20 (59%) were in patients ≥65 years of age.

Since the start of the 2014-15 season, 8,008 hospitalizations have been reported; 6,843 (86%) with influenza A. Among cases for which the subtype of influenza A was reported, 99.1% were A(H3N2). The majority of cases (70%) were ≥65 years of age (Table 6). A total of 396 ICU admissions have been reported to date: 52% (n=205) were in adults ≥65 years of age and 75% were due to influenza A. A total of 604 deaths have been reported since the start of the season: three children <5 years of age, five children 5-19 years, 47 adults 20-64 years, and 549 adults ≥65 years of age. Influenza A has been reported in 91% of deaths. Adults 65 years of age or older represent 91% of all deaths reported this season. Detailed clinical information (e.g. underlying medical conditions) is not known for these cases

* Note: Influenza-associated hospitalizations are not reported to PHAC by the following Provinces and Territory: BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by Saskatchewan. ICU admissions are not distinguished among hospital admissions reported from Ontario. Data may also include cases reported by the IMPACT and PCIRN networks. The number of new influenza-associated hospitalizations and deaths reported for the current week may include cases from Ontario that occurred in previous weeks, as a result of retrospective updates to the cumulative total. It is important to note that the hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

Table 6 – Cumulative number of hospitalizations with influenza reported by the participating provinces and territories, Canada, 2014-15

Age groups (years)	Cumulative (24 Aug. 2014 to 15 Aug. 2015)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A (UnS)	Total	# (%)
0-4	443	7	154	282	100	543 (7%)
5-19	266	0	123	143	106	372 (5%)
20-44	335	4	178	153	93	428 (5%)
45-64	748	13	368	367	226	974 (12%)
65+	4997	5	2389	2603	616	5613 (70%)
Unknown	54	0	51	3	24	78 (1%)
Total	6843	29	3263	3551	1165	8008
Percentage¹	85.5%	0.4%	47.7%	51.9%	14.5%	100.0%

¹ Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections. UnS: unsubtype: The specimen was typed as influenza A, but no result for subtyping was available.

See additional data on [Reported Influenza Hospitalizations and Deaths in Canada: 2009-10 to 2014-15](#) on the Public Health Agency of Canada website.

Emerging Respiratory Pathogens

Human Avian Influenza

Influenza A(H7N9): There have been no reported cases of human infection with avian influenza A(H7N9) since July 20, 2015. Globally to August 20, 2015, the WHO reported a total of 678 laboratory-confirmed human cases with avian influenza A(H7N9) virus, including 275 deaths. Documents related to the public health risk of influenza A(H7N9), as well as guidance for health professionals and advice for the public is updated regularly on the following websites:

[PHAC – Avian influenza A\(H7N9\)](#)

[WHO – Avian Influenza A\(H7N9\)](#)

Middle East Respiratory Syndrome Coronavirus (MERS-CoV)

Since the last FluWatch report, a number of confirmed cases of MERS-CoV have been reported in Saudi Arabia. Most of these cases have been associated with a hospital outbreak in the Riyadh Region.

There have been no new cases reported in South Korea since July 4, 2015.

Globally, from September 2012 to August 20, 2015, the WHO has reported a total of 1,432 laboratory-confirmed cases of infection with MERS-CoV, including 507 deaths.

Documents related to the public health risk of MERS-CoV, as well as guidance for health professionals and advice for the public is updated regularly on the following websites:

[PHAC – Middle East respiratory syndrome coronavirus \(MERS-CoV\)](#)

[WHO – Coronavirus infections](#)

International Influenza Reports

[World Health Organization influenza update](#)

[World Health Organization FluNet](#)

[WHO Influenza at the human-animal interface](#)

[Centers for Disease Control and Prevention seasonal influenza report](#)

[European Centre for Disease Prevention and Control - epidemiological data](#)

[South Africa Influenza surveillance report](#)

[New Zealand Public Health Surveillance](#)

[Australia Influenza Report](#)

[Pan-American Health Organization Influenza Situation Report](#)

FluWatch Definitions for the 2014-2015 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This report is available on the Public Health Agency website at the following address: <http://www.phac-aspc.gc.ca/fluwatch/index.html>.

Ce rapport est disponible dans les deux langues officielles.